

Glutaric acid, 2-fluorophenyl 2,5-dichlorophenyl ester

Inchi: InChI=1S/C17H13Cl2FO4/c18-11-8-9-12(19)15(10-11)24-17(22)7-3-6-16(21)23-14-5-2-1
InchiKey: QWHBBSSGHOWNRB-UHFFFAOYSA-N
Formula: C17H13Cl2FO4
SMILES: O=C(CCCC(=O)Oc1cc(Cl)ccc1Cl)Oc1ccccc1F
Mol. weight [g/mol]: 371.19

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|---------|----------------|
| gf | -398.32 | kJ/mol | Joback Method |
| hf | -672.75 | kJ/mol | Joback Method |
| hfus | 43.75 | kJ/mol | Joback Method |
| hvap | 86.24 | kJ/mol | Joback Method |
| log10ws | -5.87 | | Crippen Method |
| logp | 4.814 | | Crippen Method |
| mcvol | 244.000 | ml/mol | McGowan Method |
| pc | 1957.87 | kPa | Joback Method |
| rinpol | 2631.00 | | NIST Webbook |
| rinpol | 2631.00 | | NIST Webbook |
| tb | 883.37 | K | Joback Method |
| tc | 1114.35 | K | Joback Method |
| tf | 576.50 | K | Joback Method |
| vc | 0.935 | m3/kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 660.94 | J/molxK | 883.37 | Joback Method |
| cpg | 671.30 | J/molxK | 921.87 | Joback Method |
| cpg | 680.53 | J/molxK | 960.36 | Joback Method |
| cpg | 688.63 | J/molxK | 998.86 | Joback Method |
| cpg | 695.64 | J/molxK | 1037.36 | Joback Method |
| cpg | 701.57 | J/molxK | 1075.86 | Joback Method |
| cpg | 706.44 | J/molxK | 1114.35 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U392057&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinp: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/112-890-9/Glutaric-acid-2-fluorophenyl-2-5-dichlorophenyl-ester.pdf>

Generated by Cheméo on 2024-04-29 10:14:06.606473019 +0000 UTC m=+16674895.527050341.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.