

Succinic acid, 1,1,1-trifluoroprop-2-yl 3-phenylprop-2-en-1-yl ester

Inchi: InChI=1S/C16H17F3O4/c1-12(16(17,18)19)23-15(21)10-9-14(20)22-11-5-8-13-6-3-2-4-7
InchiKey: JOWXUCNFABEKIN-VMPITWQZSA-N
Formula: C16H17F3O4
SMILES: CC(OC(=O)CCC(=O)OCC=Cc1ccccc1)C(F)(F)F
Mol. weight [g/mol]: 330.30

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|---------|----------------|
| gf | -775.40 | kJ/mol | Joback Method |
| hf | -1111.78 | kJ/mol | Joback Method |
| hfus | 35.32 | kJ/mol | Joback Method |
| hvap | 67.62 | kJ/mol | Joback Method |
| log10ws | -4.14 | | Crippen Method |
| logp | 3.517 | | Crippen Method |
| mcvol | 228.430 | ml/mol | McGowan Method |
| pc | 1755.07 | kPa | Joback Method |
| rinpol | 1958.00 | | NIST Webbook |
| rinpol | 1958.00 | | NIST Webbook |
| tb | 743.04 | K | Joback Method |
| tc | 940.90 | K | Joback Method |
| tf | 424.93 | K | Joback Method |
| vc | 0.888 | m3/kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 650.97 | J/molxK | 743.04 | Joback Method |
| cpg | 664.50 | J/molxK | 776.02 | Joback Method |
| cpg | 677.10 | J/molxK | 808.99 | Joback Method |
| cpg | 688.80 | J/molxK | 841.97 | Joback Method |
| cpg | 699.66 | J/molxK | 874.95 | Joback Method |
| cpg | 709.71 | J/molxK | 907.92 | Joback Method |
| cpg | 719.01 | J/molxK | 940.90 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U391040&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307I |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/113-813-3/Succinic-acid-1-1-1-trifluoroprop-2-yl-3-phenylprop-2-en-1-yl-ester.pdf>

Generated by Cheméo on 2024-04-27 16:48:59.917942109 +0000 UTC m=+16525788.838519424.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.