

Succinic acid, hexadecyl 1-phenyl-2,2,2-trifluoroethyl ester

| | |
|-----------------------------|---|
| Inchi: | InChI=1S/C28H43F3O4/c1-2-3-4-5-6-7-8-9-10-11-12-13-14-18-23-34-25(32)21-22-26(33) |
| InchiKey: | FKPMNGPOMUMOEC-UHFFFAOYSA-N |
| Formula: | C28H43F3O4 |
| SMILES: | CCCCCCCCCCCCCCCCOC(=O)CCC(=O)OC(c1ccccc1)C(F)(F)F |
| Mol. weight [g/mol]: | 500.63 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|----------------------|----------------|
| gf | -754.58 | kJ/mol | Joback Method |
| hf | -1476.68 | kJ/mol | Joback Method |
| hfus | 66.19 | kJ/mol | Joback Method |
| hvap | 94.37 | kJ/mol | Joback Method |
| log10ws | -9.49 | | Crippen Method |
| logp | 8.638 | | Crippen Method |
| mvol | 401.810 | ml/mol | McGowan Method |
| pc | 775.48 | kPa | Joback Method |
| rinpol | 2988.00 | | NIST Webbook |
| rinpol | 2988.00 | | NIST Webbook |
| tb | 1013.44 | K | Joback Method |
| tc | 1249.15 | K | Joback Method |
| tf | 565.25 | K | Joback Method |
| vc | 1.581 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|---------|---------|-----------------|---------------|
| cpg | 1390.26 | J/mol×K | 1013.44 | Joback Method |
| cpg | 1408.58 | J/mol×K | 1052.73 | Joback Method |
| cpg | 1425.28 | J/mol×K | 1092.01 | Joback Method |
| cpg | 1440.48 | J/mol×K | 1131.30 | Joback Method |
| cpg | 1454.29 | J/mol×K | 1170.58 | Joback Method |
| cpg | 1466.83 | J/mol×K | 1209.87 | Joback Method |
| cpg | 1478.20 | J/mol×K | 1249.15 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U381583&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/115-335-2/Succinic-acid-hexadecyl-1-phenyl-2-2-2-trifluoroethyl-ester.pdf>

Generated by Cheméo on 2024-05-05 08:13:40.280565899 +0000 UTC m=+17186069.201143214.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.