

Succinic acid, 1-(2,6-difluorophenyl)ethyl hexyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C18H24F2O4/c1-3-4-5-6-12-23-16(21)10-11-17(22)24-13(2)18-14(19)8-7-9-15 |
| InchiKey: | LUZFOEHJGLRBQQ-UHFFFAOYSA-N |
| Formula: | C18H24F2O4 |
| SMILES: | CCCCCOC(=O)CCC(=O)OC(C)c1c(F)cccc1F |
| Mol. weight [g/mol]: | 342.38 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|----------------------|----------------|
| gf | -666.07 | kJ/mol | Joback Method |
| hf | -1088.36 | kJ/mol | Joback Method |
| hfus | 43.85 | kJ/mol | Joback Method |
| hvap | 75.55 | kJ/mol | Joback Method |
| log10ws | -5.31 | | Crippen Method |
| logp | 4.473 | | Crippen Method |
| mvol | 259.140 | ml/mol | McGowan Method |
| pc | 1440.25 | kPa | Joback Method |
| rinpol | 2130.00 | | NIST Webbook |
| rinpol | 2130.00 | | NIST Webbook |
| tb | 798.56 | K | Joback Method |
| tc | 991.38 | K | Joback Method |
| tf | 474.58 | K | Joback Method |
| vc | 1.014 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 776.15 | J/mol×K | 798.56 | Joback Method |
| cpg | 790.96 | J/mol×K | 830.70 | Joback Method |
| cpg | 804.79 | J/mol×K | 862.83 | Joback Method |
| cpg | 817.65 | J/mol×K | 894.97 | Joback Method |
| cpg | 829.56 | J/mol×K | 927.10 | Joback Method |
| cpg | 840.52 | J/mol×K | 959.24 | Joback Method |
| cpg | 850.54 | J/mol×K | 991.38 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U381422&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/115-717-8/Succinic-acid-1-2-6-difluorophenyl-ethyl-hexyl-ester.pdf>

Generated by Cheméo on 2024-05-01 19:36:14.655916597 +0000 UTC m=+16881423.576493918.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.