

Succinic acid, butyl 2,4,5-trifluorobenzyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C15H17F3O4/c1-2-3-6-21-14(19)4-5-15(20)22-9-10-7-12(17)13(18)8-11(10)16 |
| InchiKey: | QFVJQOCYCBUAQE-UHFFFAOYSA-N |
| Formula: | C15H17F3O4 |
| SMILES: | CCCCOC(=O)CCC(=O)OCc1cc(F)c(F)cc1F |
| Mol. weight [g/mol]: | 318.29 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|----------------------|----------------|
| gf | -893.33 | kJ/mol | Joback Method |
| hf | -1228.74 | kJ/mol | Joback Method |
| hfus | 42.29 | kJ/mol | Joback Method |
| hvap | 69.11 | kJ/mol | Joback Method |
| log10ws | -4.42 | | Crippen Method |
| logp | 3.271 | | Crippen Method |
| mvol | 218.640 | ml/mol | McGowan Method |
| pc | 1714.61 | kPa | Joback Method |
| rinpol | 1871.00 | | NIST Webbook |
| rinpol | 1871.00 | | NIST Webbook |
| tb | 734.61 | K | Joback Method |
| tc | 921.00 | K | Joback Method |
| tf | 468.88 | K | Joback Method |
| vc | 0.870 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 615.27 | J/mol×K | 734.61 | Joback Method |
| cpg | 628.26 | J/mol×K | 765.67 | Joback Method |
| cpg | 640.50 | J/mol×K | 796.74 | Joback Method |
| cpg | 651.97 | J/mol×K | 827.80 | Joback Method |
| cpg | 662.68 | J/mol×K | 858.87 | Joback Method |
| cpg | 672.63 | J/mol×K | 889.93 | Joback Method |
| cpg | 681.82 | J/mol×K | 921.00 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U382219&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinp: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/115-786-2/Succinic-acid-butyl-2-4-5-trifluorobenzyl-ester.pdf>

Generated by Cheméo on 2024-05-03 14:57:16.758738844 +0000 UTC m=+17037485.679316171.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.