

Succinic acid, 8-chlorooctyl 3-methylbut-2-yl ester

| | |
|----------------------|---|
| Inchi: | InChI=1S/C17H31ClO4/c1-14(2)15(3)22-17(20)11-10-16(19)21-13-9-7-5-4-6-8-12-18/h14 |
| InchiKey: | LFKGCNLFCDJZJX-UHFFFAOYSA-N |
| Formula: | C17H31ClO4 |
| SMILES: | CC(C)C(C)OC(=O)CCC(=O)OCCCCCCCCCl |
| Mol. weight [g/mol]: | 334.88 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -392.39 | kJ/mol | Joback Method |
| hf | -910.11 | kJ/mol | Joback Method |
| hfus | 42.51 | kJ/mol | Joback Method |
| hvap | 75.36 | kJ/mol | Joback Method |
| log10ws | -4.69 | | Crippen Method |
| logp | 4.477 | | Crippen Method |
| mvol | 277.510 | ml/mol | McGowan Method |
| pc | 1294.86 | kPa | Joback Method |
| rinpol | 2278.00 | | NIST Webbook |
| rinpol | 2278.00 | | NIST Webbook |
| tb | 777.49 | K | Joback Method |
| tc | 963.26 | K | Joback Method |
| tf | 425.59 | K | Joback Method |
| vc | 1.073 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 823.50 | J/molxK | 777.49 | Joback Method |
| cpg | 896.14 | J/molxK | 932.30 | Joback Method |
| cpg | 883.46 | J/molxK | 901.34 | Joback Method |
| cpg | 869.87 | J/molxK | 870.38 | Joback Method |
| cpg | 855.35 | J/molxK | 839.41 | Joback Method |
| cpg | 839.90 | J/molxK | 808.45 | Joback Method |
| cpg | 907.93 | J/molxK | 963.26 | Joback Method |
| dvisc | 0.0000598 | Paxs | 777.49 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0000811 | Paxs | 718.84 | Joback Method |
| dvisc | 0.0001159 | Paxs | 660.19 | Joback Method |
| dvisc | 0.0001778 | Paxs | 601.54 | Joback Method |
| dvisc | 0.0002990 | Paxs | 542.89 | Joback Method |
| dvisc | 0.0005702 | Paxs | 484.24 | Joback Method |
| dvisc | 0.0012997 | Paxs | 425.59 | Joback Method |

Sources

| | |
|------------------------|---|
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U390589&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/119-068-5/Succinic-acid-8-chlorooctyl-3-methylbut-2-yl-ester.pdf>

Generated by Cheméo on 2024-04-29 10:04:38.819362172 +0000 UTC m=+16674327.739939494.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.