

L-2-Aminobutyric acid, N-dimethylaminomethylene-, ethyl ester

Inchi: InChI=1S/C9H18N2O2/c1-5-8(9(12)13-6-2)10-7-11(3)4/h7-8H,5-6H2,1-4H3
InchiKey: ODSCIKNBQRNJGD-UHFFFAOYSA-N
Formula: C9H18N2O2
SMILES: CCOC(=O)C(CC)N=CN(C)C
Mol. weight [g/mol]: 186.25

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|--------|----------------|
| hf | -329.42 | kJ/mol | Joback Method |
| hvap | 49.75 | kJ/mol | Joback Method |
| log10ws | -0.79 | | Crippen Method |
| logp | 0.918 | | Crippen Method |
| mcvol | 160.770 | ml/mol | McGowan Method |
| pc | 2193.84 | kPa | Joback Method |
| rinpol | 1301.00 | | NIST Webbook |
| rinpol | 1301.00 | | NIST Webbook |
| tb | 570.29 | K | Joback Method |
| tc | 762.15 | K | Joback Method |

Sources

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>
Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws
Joback Method: https://en.wikipedia.org/wiki/Joback_method
McGowan Method: <http://link.springer.com/article/10.1007/BF02311772>
NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=U375526&Units=SI>

Legend

hf: Enthalpy of formation at standard conditions
hvap: Enthalpy of vaporization at standard conditions

| | |
|-----------------|-------------------------------------|
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |

Latest version available from:

<https://www.cheméo.com/cid/119-785-9/L-2-Aminobutyric-acid-N-dimethylaminomethylene-ethyl-ester.pdf>

Generated by Cheméo on 2024-05-03 02:18:46.786782089 +0000 UTC m=+16991975.707359403.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.