

C-(2,3-Dihydroxypropyl)glyceraric acid, TMS

Inchi: InChI=1S/C21H50O7Si5/c1-29(2,3)24-17-18(25-30(4,5)6)16-21(28-33(13,14)15,19(22)20)26
InchiKey: ZCSYALQZRWXVJO-UHFFFAOYSA-N
Formula: C₂₁H₅₀O₇Si₅
SMILES: C[Si](C)(C)OCC(CC(O[Si](C)(C)C)(C(=O)O[Si](C)(C)C)C(=O)O[Si](C)(C)C)O[Si](C)(C)C
Mol. weight [g/mol]: 555.04

Physical Properties

Property code	Value	Unit	Source
log10ws	5.92		Crippen Method
logp	5.795		Crippen Method
rinpol	1949.00		NIST Webbook
rinpol	1949.00		NIST Webbook

Sources

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>
Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws
NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=R101609&Units=SI>

Legend

log10ws: Log10 of Water solubility in mol/l
logp: Octanol/Water partition coefficient
rinpol: Non-polar retention indices

Latest version available from:

<https://www.chemeo.com/cid/123-497-4/C-2-3-Dihydroxypropyl-glyceraric-acid-TMS.pdf>

Generated by Cheméo on 2024-04-30 07:50:16.270178778 +0000 UTC m=+16752665.190756095.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.