

Succinic acid, 2-methylpent-3-yl 2-chlorophenyl ester

| | |
|-----------------------------|---|
| Inchi: | InChI=1S/C16H21ClO4/c1-4-13(11(2)3)20-15(18)9-10-16(19)21-14-8-6-5-7-12(14)17/h5- |
| InchiKey: | WBPXLYAGSOWQGW-UHFFFAOYSA-N |
| Formula: | C16H21ClO4 |
| SMILES: | CCC(OC(=O)CCC(=O)Oc1ccccc1Cl)C(C)C |
| Mol. weight [g/mol]: | 312.79 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -298.03 | kJ/mol | Joback Method |
| hf | -664.41 | kJ/mol | Joback Method |
| hfus | 33.57 | kJ/mol | Joback Method |
| hvap | 76.07 | kJ/mol | Joback Method |
| log10ws | -4.55 | | Crippen Method |
| logp | 4.003 | | Crippen Method |
| mvol | 239.660 | ml/mol | McGowan Method |
| pc | 1784.86 | kPa | Joback Method |
| rinpol | 2093.00 | | NIST Webbook |
| rinpol | 2093.00 | | NIST Webbook |
| tb | 786.27 | K | Joback Method |
| tc | 997.58 | K | Joback Method |
| tf | 453.26 | K | Joback Method |
| vc | 0.908 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 676.33 | J/molxK | 786.27 | Joback Method |
| cpg | 690.72 | J/molxK | 821.49 | Joback Method |
| cpg | 704.05 | J/molxK | 856.71 | Joback Method |
| cpg | 716.33 | J/molxK | 891.92 | Joback Method |
| cpg | 727.57 | J/molxK | 927.14 | Joback Method |
| cpg | 737.79 | J/molxK | 962.36 | Joback Method |
| cpg | 747.01 | J/molxK | 997.58 | Joback Method |
| dvisc | 0.0009423 | Paxs | 453.26 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0004823 | Paxs | 508.76 | Joback Method |
| dvisc | 0.0002816 | Paxs | 564.26 | Joback Method |
| dvisc | 0.0001811 | Paxs | 619.76 | Joback Method |
| dvisc | 0.0001252 | Paxs | 675.27 | Joback Method |
| dvisc | 0.0000916 | Paxs | 730.77 | Joback Method |
| dvisc | 0.0000700 | Paxs | 786.27 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U389797&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|----------------------------|---|
| cp_g: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log₁₀ws: | Log ₁₀ of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/125-019-2/Succinic-acid-2-methylpent-3-yl-2-chlorophenyl-ester.pdf>

Generated by Cheméo on 2024-05-01 22:55:15.504387751 +0000 UTC m=+16893364.424965066.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.