

Sebacic acid, 1-phenyl-2,2,2-trifluoromethylethyl propyl

Inchi:
ester

InChI=1S/C21H29F3O4/c1-2-16-27-18(25)14-10-5-3-4-6-11-15-19(26)28-20(21(22,23)24

InchiKey:

RVVXQSQGXWWEPS-UHFFFAOYSA-N

Formula:

C21H29F3O4

SMILES:

CCCOC(=O)CCCCCCCC(=O)OC(c1ccccc1)C(F)(F)F

Mol. weight [g/mol]:

402.45

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|----------------------|----------------|
| gf | -813.52 | kJ/mol | Joback Method |
| hf | -1332.20 | kJ/mol | Joback Method |
| hfus | 48.06 | kJ/mol | Joback Method |
| hvap | 78.79 | kJ/mol | Joback Method |
| log10ws | -6.56 | | Crippen Method |
| logp | 5.907 | | Crippen Method |
| mvol | 303.180 | ml/mol | McGowan Method |
| pc | 1173.63 | kPa | Joback Method |
| rinpol | 2261.00 | | NIST Webbook |
| rinpol | 2261.00 | | NIST Webbook |
| tb | 853.28 | K | Joback Method |
| tc | 1048.99 | K | Joback Method |
| tf | 486.36 | K | Joback Method |
| vc | 1.188 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|---------|---------|-----------------|---------------|
| cpg | 960.99 | J/mol×K | 853.28 | Joback Method |
| cpg | 976.51 | J/mol×K | 885.90 | Joback Method |
| cpg | 990.94 | J/mol×K | 918.52 | Joback Method |
| cpg | 1004.31 | J/mol×K | 951.13 | Joback Method |
| cpg | 1016.67 | J/mol×K | 983.75 | Joback Method |
| cpg | 1028.08 | J/mol×K | 1016.37 | Joback Method |
| cpg | 1038.57 | J/mol×K | 1048.99 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U416800&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307I |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinp: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/125-264-0/Sebacic-acid-1-phenyl-2-2-2-trifluoromethylethyl-propyl-ester.pdf>

Generated by Cheméo on 2024-04-29 20:41:31.922365544 +0000 UTC m=+16712540.842942900.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.