

2,3-dimethyl-4,5-dithiaoctane

| | |
|-----------------------------|--|
| Other names: | Propyl-sec.isoamyldisulfide |
| Inchi: | InChI=1S/C8H18S2/c1-5-6-9-10-8(4)7(2)3/h7-8H,5-6H2,1-4H3 |
| InchiKey: | NWRHSMBBPGLJSE-UHFFFAOYSA-N |
| Formula: | C8H18S2 |
| SMILES: | CCCSSC(C)C(C)C |
| Mol. weight [g/mol]: | 178.36 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 77.84 | kJ/mol | Joback Method |
| hf | -135.27 | kJ/mol | Joback Method |
| hfus | 17.69 | kJ/mol | Joback Method |
| hvap | 46.26 | kJ/mol | Joback Method |
| log10ws | -3.80 | | Crippen Method |
| logp | 3.822 | | Crippen Method |
| mcpol | 156.280 | ml/mol | McGowan Method |
| pc | 2668.02 | kPa | Joback Method |
| rinpol | 1237.00 | | NIST Webbook |
| rinpol | 1235.00 | | NIST Webbook |
| rinpol | 1237.00 | | NIST Webbook |
| rinpol | 1235.00 | | NIST Webbook |
| tb | 519.12 | K | Joback Method |
| tc | 734.48 | K | Joback Method |
| tf | 218.72 | K | Joback Method |
| vc | 0.580 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 341.26 | J/mol×K | 519.12 | Joback Method |
| cpg | 356.56 | J/mol×K | 555.01 | Joback Method |
| cpg | 371.10 | J/mol×K | 590.91 | Joback Method |
| cpg | 384.91 | J/mol×K | 626.80 | Joback Method |
| cpg | 397.97 | J/mol×K | 662.70 | Joback Method |

| | | | | |
|-----|--------|---------|--------|---------------|
| cpg | 410.31 | J/mol×K | 698.59 | Joback Method |
| cpg | 421.93 | J/mol×K | 734.48 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=R155507&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvac: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/15-092-3/2-3-dimethyl-4-5-dithiaoctane.pdf>

Generated by Cheméo on 2024-04-28 04:18:30.418765186 +0000 UTC m=+16567159.339342498.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.