

Benzoic acid, 3-(isopropylthio)-, isopropyl ester

| | |
|----------------------|--|
| Inchi: | InChI=1S/C13H18O2S/c1-9(2)15-13(14)11-6-5-7-12(8-11)16-10(3)4/h5-10H,1-4H3 |
| InchiKey: | PFNXMWHIBDDMBH-UHFFFAOYSA-N |
| Formula: | C13H18O2S |
| SMILES: | CC(C)OC(=O)c1cccc(SC(C)C)c1 |
| Mol. weight [g/mol]: | 238.35 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -44.32 | kJ/mol | Joback Method |
| hf | -300.08 | kJ/mol | Joback Method |
| hfus | 22.95 | kJ/mol | Joback Method |
| hvap | 62.67 | kJ/mol | Joback Method |
| log10ws | -4.36 | | Crippen Method |
| logp | 3.752 | | Crippen Method |
| mcvol | 194.060 | ml/mol | McGowan Method |
| pc | 2331.52 | kPa | Joback Method |
| rinpol | 1704.00 | | NIST Webbook |
| rinpol | 1704.00 | | NIST Webbook |
| tb | 672.69 | K | Joback Method |
| tc | 901.16 | K | Joback Method |
| tf | 351.77 | K | Joback Method |
| vc | 0.722 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 500.34 | J/mol×K | 672.69 | Joback Method |
| cpg | 516.33 | J/mol×K | 710.77 | Joback Method |
| cpg | 531.24 | J/mol×K | 748.85 | Joback Method |
| cpg | 545.08 | J/mol×K | 786.92 | Joback Method |
| cpg | 557.87 | J/mol×K | 825.00 | Joback Method |
| cpg | 569.62 | J/mol×K | 863.08 | Joback Method |
| cpg | 580.37 | J/mol×K | 901.16 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U375401&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/16-136-3/Benzoic-acid-3-isopropylthio-isopropyl-ester.pdf>

Generated by Cheméo on 2024-04-18 23:56:06.786405141 +0000 UTC m=+15773815.706982463.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.