1-Heptyne

Other names:	AMYLACETYLENE
	hept-1-yne
	n-C5H11C«equiv»CH
	n-C5H11CÂ«equivÂ»CH
Inchi:	InChI=1S/C7H12/c1-3-5-7-6-4-2/h1H,4-7H2,2H3
InchiKey:	YVXHZKKCZYLQOP-UHFFFAOYSA-N
Formula:	C7H12
SMILES:	C#CCCCCC
Mol. weight [g/mol]:	96.17
CAS:	628-71-7

Physical Properties

Property code	Value	Unit	Source
af	0.2930		KDB
chl	-4570.60	kJ/mol	NIST Webbook
gf	231.13	kJ/mol	Joback Method
hcg	4542.99	kJ/mol	KDB
hcn	4279.395	kJ/mol	KDB
hf	101.70	kJ/mol	NIST Webbook
hf	103.80 ± 2.60	kJ/mol	NIST Webbook
hfl	101.10 ± 4.00	kJ/mol	NIST Webbook
hfl	-62.80	kJ/mol	NIST Webbook
hfus	16.86	kJ/mol	Joback Method
hvap	31.03	kJ/mol	Joback Method
ie	10.04 ± 0.01	eV	NIST Webbook
log10ws	-3.01		Estimated Solubility Method
log10ws	-3.01		Aqueous Solubility Prediction Method
logp	2.200		Crippen Method
mcvol	100.890	ml/mol	McGowan Method
pc	3300.00	kPa	KDB
rinpol	686.00		NIST Webbook
rinpol	686.00		NIST Webbook
rinpol	712.00		NIST Webbook
rinpol	686.00		NIST Webbook
rinpol	686.00		NIST Webbook
rinpol	687.00		NIST Webbook

rinpol	686.00		NIST Webbook
rinpol	687.30		NIST Webbook
rinpol	688.20		NIST Webbook
rinpol	689.00		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	711.00		NIST Webbook
rinpol	685.30		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	686.00		NIST Webbook
rinpol	684.00		NIST Webbook
rinpol	687.10		NIST Webbook
rinpol	687.00		NIST Webbook
rinpol	686.00		NIST Webbook
ripol	947.00		NIST Webbook
ripol	938.00		NIST Webbook
ripol	929.00		NIST Webbook
ripol	934.00		NIST Webbook
tb	373.15 ± 3.00	K	NIST Webbook
tb	372.15 ± 2.00	K	NIST Webbook
tb	372.15 ± 2.00	K	NIST Webbook
tb	372.15 ± 2.00	K	NIST Webbook
tb	372.65 ± 2.00	K	NIST Webbook
tb	372.99 ± 0.50	K	NIST Webbook
tb	379.15 ± 5.00	K	NIST Webbook
tb	372.90	K	KDB
tb	372.50 ± 0.50	K	NIST Webbook
tb	372.90	K	NIST Webbook
tb	372.90 ± 1.50	K	NIST Webbook
tb	371.15 ± 1.50	K	NIST Webbook
tb	374.00 ± 2.00	K	NIST Webbook
tb	371.65 ± 2.00	K	NIST Webbook
tb	371.15 ± 1.50	K	NIST Webbook
tb	372.65 ± 0.70	K	NIST Webbook
tb	372.89 ± 0.30	K	NIST Webbook
tb	372.89 ± 0.40	K	NIST Webbook
tb	372.93 ± 0.20	K	NIST Webbook
tb	372.15 ± 1.50	K	NIST Webbook
tb	360.15	K	NIST Webbook
tb	371.75 ± 1.50	K	NIST Webbook
tb	372.15 ± 1.50	K	NIST Webbook
tb	371.15 ± 0.50	K	NIST Webbook

tb	371.65 ± 1.50	K	NIST Webbook
tb	371.65 ± 1.00	K	NIST Webbook
tb	372.65 ± 1.50	K	NIST Webbook
tb	373.75 ± 1.50	K	NIST Webbook
tb	372.65 ± 1.50	K	NIST Webbook
tb	371.15 ± 1.50	K	NIST Webbook
tb	373.15 ± 1.50	K	NIST Webbook
tb	373.15 ± 1.50	K	NIST Webbook
tb	371.65 ± 1.00	K	NIST Webbook
tb	373.15 ± 1.50	K	NIST Webbook
tc	559.70	K	KDB
tf	192.00	K	KDB
tf	192.10 ± 0.50	K	NIST Webbook
tf	192.22 ± 0.10	K	NIST Webbook
tf	192.15 ± 1.50	K	NIST Webbook
vc	0.390	m3/kmol	KDB
zc	0.2762040		KDB

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	210.88	J/mol×K	466.49	Joback Method
cpg	219.59	J/mol×K	495.70	Joback Method
cpg	172.06	J/mol×K	349.68	Joback Method
cpg	182.39	J/mol×K	378.88	Joback Method
cpg	192.30	J/mol×K	408.09	Joback Method
cpg	201.79	J/mol×K	437.29	Joback Method
cpg	227.93	J/mol×K	524.90	Joback Method
hvapt	37.90	kJ/mol	354.50	NIST Webbook
rfi	1.40610		298.15	KDB

Correlations

Information	Value
Property code	pvap
Equation	ln(Pvp) = A + B/(T + C)
Coeff. A	1.38643e+01
Coeff. B	-2.91233e+03

Coeff. C	-5.79460e+01
Temperature range (K), min.	272.46
Temperature range (K), max.	398.46

Information	Value
Property code	pvap
Equation	ln(Pvp) = A + B/T + C*ln(T) + D*T^2
Coeff. A	8.20864e+01
Coeff. B	-6.66117e+03
Coeff. C	-1.02999e+01
Coeff. D	9.86822e-06
Temperature range (K), min.	287.15
Temperature range (K), max.	559.69

Sources

Activity coefficients at infinite dilution for solutes in the trioctylmethylammonium	https://www.doi.org/10.1016/j.jct.2009.08.012
4-1 alkylpyridinium-based ionic liquid in the separation processes:	https://www.doi.org/10.1016/j.jct.2016.01.017
Bis(trifluoromethylsulfonyl)imide ionic liquid using gas liquid	https://www.doi.org/10.1016/j.jct.2014.04.024
Thermodynamics and activity coefficients at infinite dilution for	https://www.doi.org/10.1016/j.fluid.2014.06.021
Experimental and theoretical study of organic solutes, water and diols in the	https://www.doi.org/10.1016/j.jct.2008.01.004
Interaction between organic	https://www.doi.org/10.1016/j.jct.2012.05.017
Measurements of activity coefficients at infinite dilution of aromatic and	https://www.doi.org/10.1021/acs.jced.8b00080
Bis(trifluoromethylsulfonyl)imide:	https://www.doi.org/10.1016/j.jct.2015.05.014
Activity coefficients at infinite dilution of organic solutes and water in the new ionic liquid	https://www.doi.org/10.1016/j.fluid.2017.06.001
Activity coefficients at infinite dilution and physicochemical properties for	https://www.doi.org/10.1016/j.jct.2016.06.028
Determination of Activity Coefficients at Infinite Dilution of Solutes in the	https://www.doi.org/10.1016/j.jct.2013.08.030
Activity coefficients at infinite dilution for various solutes in	https://www.doi.org/10.1016/j.jct.2013.05.030
Activity coefficients at infinite dilution of organic solutes in the ionic liquid	https://www.doi.org/10.1021/je500050p
Activity Coefficients at Infinite Dilution Determination of Activity Coefficients at Infinite Dilution of Solutes in the	https://www.doi.org/10.1021/je800105r
Activity coefficients at infinite dilution of organic solutes in the ionic liquid of water/butan-1-ol based on activity coefficients at infinite	https://www.doi.org/10.1016/j.jct.2017.10.003
Thermodynamics and limiting activity coefficients of organic solutes in the ionic liquid	https://www.doi.org/10.1016/j.jct.2016.08.008
Activity Coefficients at Infinite Dilution of Organic Solutes in the Ionic Liquid	https://www.doi.org/10.1016/j.jct.2011.04.018
Estimated Solubility Methods:	http://pubs.acs.org/doi/suppl/10.1021/ci034243x/suppl_file/ci034243xsi20040112_053635.txt
Evaluation of the performance of	https://www.doi.org/10.1021/je201129y
Trigeminal Tricationic Ionic Liquids for	https://www.doi.org/10.1021/je030187k
Determination of Activity Coefficients at Infinite Dilution of Solutes in the Ionic Liquid	https://www.doi.org/10.1016/j.jct.2010.01.004
measurements of activity coefficients at infinite dilution	https://www.doi.org/10.1021/je060033f
Thermodynamic Properties of Mixtures Containing Ionic Liquids: Activity	https://www.doi.org/10.1016/j.jct.2015.05.022
Activity coefficients at infinite dilution,	https://www.doi.org/10.1016/j.jct.2011.11.025
Organic Compounds in 1-Propyl-	
Boronic Acid 3-Pyridylmidazolium	
Boronic acid:	
Activity coefficients of organic solutes and water in the ionic liquid	
1-Propyl-3-pyridylmidazolium Bromide	
1-ethyl-3-methylimidazolium phosphate:	
tetracyanoborate:	

The use of ionic liquids for separation of binary hydrocarbons mixtures based on gamma infinity data measurements:	https://www.doi.org/10.1016/j.jct.2018.07.024
Thermodynamics and activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid [illegible]	https://www.doi.org/10.1016/j.jct.2011.06.007
Thermodynamic study of molecular interaction selectivity in separation [illegible]	https://www.doi.org/10.1016/j.jct.2018.02.014
Activity coefficients at infinite dilution of organic solutes in the ionic liquid [illegible]	https://www.doi.org/10.1016/j.jct.2009.12.004
Measurements of activity coefficients at infinite dilution in [illegible] hexafluorophosphate using gas-liquid chromatography at T = (313.15, [illegible]	https://www.doi.org/10.1016/j.jct.2007.01.004
Activity coefficients at infinite dilution of organic solutes in the ionic liquid [illegible]	https://www.doi.org/10.1016/j.jct.2010.12.019
Activity coefficients at infinite dilution measurements for organic solutes in [illegible] chromatography at T = (313.15, 323.15, [illegible]	https://www.doi.org/10.1016/j.jct.2005.01.015
Activity coefficients at infinite dilution measurements for organic solutes and water in [illegible]	https://www.doi.org/10.1016/j.fluid.2009.08.017
Activity Coefficients at Infinite Dilution of Organic Solutes and Water in [illegible]	https://www.doi.org/10.1021/acs.jced.5b00980
Separation of [illegible] aromatic hydrocarbons and sulfur compounds [illegible]	https://www.doi.org/10.1016/j.jct.2016.07.017
[illegible]	https://www.doi.org/10.1016/j.fluid.2006.07.015
[illegible] hydrocarbons in C78H158 branched [illegible]	http://pubs.acs.org/doi/abs/10.1021/ci990307l
Abraham model linear free energy relationships for describing the partitioning and solubility behavior of [illegible]	https://www.doi.org/10.1016/j.fluid.2016.10.009
[illegible]	https://www.doi.org/10.1016/j.jct.2015.08.017
Measurement of activity coefficients at infinite dilution of organic solutes in [illegible]	https://www.doi.org/10.1016/j.jct.2013.10.017
Ammonium ionic liquids in extraction [illegible]	https://www.doi.org/10.1016/j.fluid.2018.09.024
Activity coefficients at infinite dilution [illegible]	https://www.doi.org/10.1016/j.jct.2012.08.016
[illegible]	https://www.cheric.org/research/kdb/hcprop/showprop.php?cmpid=414
Activity Coefficients at Infinite Dilution for Organic Compounds Dissolved in [illegible]	https://www.doi.org/10.1021/je300692s
Activity coefficients at infinite dilution of organic solutes in [illegible]	https://www.doi.org/10.1016/j.jct.2013.05.011
Measurement of activity coefficients at infinite dilution using polar and non-polar solutes in the ionic liquid [illegible]	https://www.doi.org/10.1016/j.jct.2005.07.003
[illegible] activity coefficients at infinite dilution for organic solutes in [illegible]	https://www.doi.org/10.1016/j.fluid.2018.11.011
Activity Coefficients at Infinite Dilution [illegible]	https://www.doi.org/10.1021/je800658v
Separation [illegible] based on limiting activity coefficients data using [illegible]	https://www.doi.org/10.1016/j.fluid.2017.12.029
Activity coefficients at infinite dilution measurements for organic solutes in [illegible]	https://www.doi.org/10.1016/j.jct.2008.04.002
Measurements of activity coefficients at infinite dilution for organic solutes in [illegible]	https://www.doi.org/10.1016/j.jct.2012.03.005
Thermodynamics and activity coefficients at infinite dilution for [illegible]	https://www.doi.org/10.1016/j.fluid.2018.06.013
[illegible] activity coefficients at Infinite Dilution of Solutes in the [illegible]	https://www.doi.org/10.1021/je0500375
Tetramethylammonium chloride + glycerol deep eutectic solvent [illegible]	https://www.doi.org/10.1016/j.fluid.2018.06.003
Activity coefficients at infinite dilution of organic solutes in the ionic liquid [illegible]	https://www.doi.org/10.1016/j.jct.2009.07.010
Activity coefficients at infinite dilution [illegible]	https://www.doi.org/10.1016/j.jct.2018.01.003
Activity Coefficients at Infinite Dilution [illegible]	https://www.doi.org/10.1021/acs.jced.8b00600
Activity coefficients at infinite dilution [illegible]	https://www.doi.org/10.1016/j.jct.2012.03.015
Measurements of activity coefficients at infinite dilution for organic solutes [illegible]	https://www.doi.org/10.1016/j.jct.2013.02.004
Assessment of Pyrrolidinium-Based [illegible]	https://www.doi.org/10.1021/acs.jced.9b00341
[illegible] Binary Mixtures Based on Activity Coefficients at Infinite Dilution: [illegible]	https://www.doi.org/10.1016/j.jct.2017.11.017
Activity coefficients at infinite dilution measurements for organic solutes and water in [illegible]	https://www.doi.org/10.1016/j.jct.2009.06.011
Activity coefficients at infinite dilution of organic solvents and water in [illegible]	https://www.doi.org/10.1016/j.fluid.2016.02.004
Study of interaction between organic compounds and [illegible]	https://www.doi.org/10.1016/j.fluid.2014.11.020
Relation between characteristic molecular volume and hydrophobicity of nonpolar molecules:	https://www.doi.org/10.1016/j.jct.2010.04.011
Measurements of activity coefficient at infinite dilution for organic solutes in tetramethylammonium chloride + ethylene glycol deep eutectic solvent using gas-liquid chromatography:	https://www.doi.org/10.1016/j.fluid.2018.01.019
Aqueous Solubility Prediction Method:	http://onschallenge.wikispaces.com/file/view/AqueousDataset002.xlsx/351826032/AqueousDa
Measurements of activity coefficients at infinite dilution of aliphatic and aromatic hydrocarbons, alcohols, thiophene, tetrahydrofuran, MTBE, and water in ionic liquid [BMIM][SCN] using GLC:	https://www.doi.org/10.1016/j.jct.2008.12.018

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 1-hexyl-3-methylimidazolium bis(trifluoromethylsulfonyl)-imide using g.l.c. at T = (298.15, 313.15, and 333.15) K: https://www.doi.org/10.1016/j.fluid.2008.10.008

Physicochemical properties and activity coefficients at infinite dilution measurements for organic solutes in 1-butyl-1-methylpyrrolidinium tetracyanoborate ionic liquid: https://www.doi.org/10.1016/j.jct.2012.09.033

Partition Coefficients of Organic Compounds in New Imidazolium and Tetraalkylammonium Based Ionic Liquids Using Inverse Gas Chromatography: https://www.doi.org/10.1021/je200637v

Activity Coefficients at Infinite Dilution Measurements for Organic Solutes and Water in the Ionic Liquid 1-Ethyl-3-methylimidazolium Ethyl Sulfate: https://www.doi.org/10.1021/je900890u

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-hexyl-3-methylimidazolium tetracyanoborate: https://www.doi.org/10.1016/j.jct.2011.02.012

Activity coefficients at infinite dilution measurements of organic solutes and water in the ionic liquid 1-hexyl-3-methylimidazolium dicyanamide: https://www.doi.org/10.1016/j.jct.2010.05.017

Experimental and theoretical study on infinite dilution activity coefficients of organic solutes in 1-hexyl-3-methylimidazolium hexafluorophosphate from Gas-Liquid Chromatography: https://www.doi.org/10.1016/j.jct.2013.01.005

Activity Coefficients at Infinite Dilution of Organic Solutes in 1-Hexyl-3-methylimidazolium Bis(trifluoromethylsulfonyl)imide Using Inverse Gas Chromatography: https://www.doi.org/10.1021/je0256481

Activity coefficients at infinite dilution and physicochemical properties for organic solutes and water in the ionic liquid 1-(2-hydroxyethyl)-3-methylimidazolium dicyanamide: https://www.doi.org/10.1021/je800754w

Activity coefficients at infinite dilution and physicochemical properties for organic solutes and water in the ionic liquid 1-(2-hydroxyethyl)-3-methylimidazolium bis(trifluoromethylsulfonyl)-imide: http://link.springer.com/article/10.1007/BF02311772

Activity coefficients at infinite dilution measurements of organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2013.05.008

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 4-(2-methoxyethyl)-4-methylmorpholinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2013.01.007

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpiperidinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2013.09.007

Activity Coefficients at Infinite Dilution of Organic Compounds in Four New Tetraalkylammonium Based Ionic Liquids Using Inverse Gas Chromatography: https://www.doi.org/10.1021/je900838a

Joback Method: https://en.wikipedia.org/wiki/Joback_method

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 1-hexyl-3-methyl-imidazolium bis(trifluoromethylsulfonyl)-imide using g.l.c. at T = (298.15, 313.15, and 333.15) K: https://www.doi.org/10.1016/j.jct.2005.03.014

The Yaws Handbook of Vapor Pressure: https://www.sciencedirect.com/book/9780128029992/the-yaws-handbook-of-vapor-pressure

KDB: https://www.cheric.org/files/research/kdb/mol/mol414.mol

Partition Coefficients of Organic Compounds in New Imidazolium and Tetraalkylammonium Based Ionic Liquids Using Inverse Gas Chromatography: https://www.doi.org/10.1021/je9003178

Gas–liquid chromatography measurements of activity coefficients at infinite dilution of hydrocarbons and alcohols in 1-alkyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide and 1-alkyl-1-methylpyrrolidinium bis(trifluoromethylsulfonyl)imide ionic liquids: https://www.doi.org/10.1016/j.fluid.2010.08.016

Activity Coefficients at Infinite Dilution of Organic Compounds in Four New Tetraalkylammonium Based Ionic Liquids Using Inverse Gas Chromatography: https://www.doi.org/10.1016/j.jct.2013.02.006

NIST Webbook: http://webbook.nist.gov/cgi/cbook.cgi?ID=C628717&Units=SI

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium dicyanamide: https://www.doi.org/10.1016/j.jct.2008.12.005

Activity coefficients at infinite dilution and physicochemical properties for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpiperidinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2012.01.019

Measurements of activity coefficients at infinite dilution for organic solutes in the ionic liquid 4-(2-methoxyethyl)-4-methylmorpholinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2013.07.004

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium tricyanomethanide: https://www.doi.org/10.1016/j.fluid.2009.01.011

Activity coefficients at infinite dilution of organic solutes in the ionic liquid 1-butyl-1-methylpyrrolidinium dicyanamide: https://www.doi.org/10.1016/j.jct.2010.02.006

Activity coefficient measurements at infinite dilution and physicochemical properties for organic solutes and water in the ionic liquid 1-butyl-1-methylpyrrolidinium dicyanamide: https://www.doi.org/10.1016/j.jct.2011.11.021

Activity coefficients at infinite dilution measurements for hydrocarbons in furfuryl alcohol at a temperature of 298.15 K, determined by gas–liquid chromatography: https://www.doi.org/10.1016/j.jct.2004.03.001

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 4-(2-methoxyethyl)-4-methylmorpholinium bis(trifluoromethylsulfonyl)-amide: https://www.doi.org/10.1016/j.fluid.2019.03.023

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid trihexyltetradecylphosphonium dicyanamide ionic liquid for various applications: https://www.doi.org/10.1016/j.jct.2014.12.027

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid trihexyltetradecylphosphonium tetracyanoborate: https://www.doi.org/10.1016/j.fluid.2005.04.021

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium tricyanomethanide: https://www.doi.org/10.1021/je0498107

Activity coefficients at infinite dilution of organic solutes in the ionic liquid trihexyltetradecylphosphonium tetracyanoborate using GLC: https://www.doi.org/10.1016/j.jct.2017.03.004

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium dicyanamide: https://www.doi.org/10.1016/j.jct.2010.12.005

Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-(2-methoxyethyl)-1-methylpyrrolidinium bis(trifluoromethylsulfonyl)-imide: https://www.doi.org/10.1016/j.jct.2012.01.004

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 1-butyl-1-methylpyrrolidinium tetracyanoborate using gas–liquid chromatography at T = (313.15, 333.15, and 353.15) K: https://www.doi.org/10.1021/je0602925

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 1-hexyl-3-methylimidazolium tetracyanoborate using gas–liquid chromatography at T = (313.15, 333.15, and 353.15) K: https://www.doi.org/10.1021/je100410k

Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid [P4,4,4,4][NTf2] in bis-butanol extraction on investigation using gas–liquid chromatography: https://www.doi.org/10.1016/j.fluid.2018.07.028

Activity coefficients at infinite dilution of organic solutes in 1-butyl-3-methylimidazolium dibutylphosphate ionic liquid using gas–liquid chromatography: https://www.doi.org/10.1016/j.jct.2015.02.023

Activity coefficients at infinite dilution measurements for organic solutes in methylphosphonate based ionic liquids using gas–liquid chromatography: https://www.doi.org/10.1016/j.jct.2015.02.024

Measurements of activity coefficients at infinite dilution of organic compounds and water in isoquinolinium based ionic liquid 4-(3-hydroxypropyl)-4-methylmorpholinium bis(trifluoromethylsulfonyl)-amide [C8iQuin][NTf2] using GLC: https://www.doi.org/10.1016/j.jct.2010.10.026

Activity Coefficients at Infinite Dilution Measurements for Organic Solutes and Water in the Ionic Liquid 1-Ethyl-3-methylimidazolium Ethylsulfate	https://www.doi.org/10.1021/je1000582
The determination of activity coefficients at infinite dilution using inverse gas chromatography.	https://www.doi.org/10.1016/j.jct.2003.09.011
Measurements of Activity Coefficients at Infinite Dilution of Organic Solutes and Water in the Ionic Liquid 1-Butyl-1-methylpiperydinium Trifluoromethanesulfonate.	https://www.doi.org/10.1021/je101008y
Activity Coefficients at Infinite Dilution of Organic Solutes in 1-Ethyl-3-methylimidazolium Tetracyanoborate Ionic Liquid Measured by Inverse Gas Chromatography.	https://www.doi.org/10.1021/je9008443
Activity coefficients at infinite dilution of organic solutes in 1-alkyl-1-methylpyrrolidinium bis(trifluoromethylsulfonyl)imide ionic liquids by gas–liquid chromatography: 2, 4, 5) using gas chromatography.	https://www.doi.org/10.1016/j.jct.2010.06.009
Activity Coefficients at Infinite Dilution for Organic Solutes Dissolved in Three 1-Alkyl-1-methylpyrrolidinium Bis(trifluoromethylsulfonyl)imide Ionic Liquids Bearing Short Linear Alkyl Side Chains of Three to Five Carbons:	https://www.doi.org/10.1021/je4001894

Legend

af:	Acentric Factor
chl:	Standard liquid enthalpy of combustion
cpg:	Ideal gas heat capacity
gf:	Standard Gibbs free energy of formation
hcg:	Heat of Combustion, Gross form
hcn:	Heat of Combustion, Net Form
hf:	Enthalpy of formation at standard conditions
hfl:	Liquid phase enthalpy of formation at standard conditions
hfus:	Enthalpy of fusion at standard conditions
hvap:	Enthalpy of vaporization at standard conditions
hvapt:	Enthalpy of vaporization at a given temperature
ie:	Ionization energy
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
pc:	Critical Pressure
pvap:	Vapor pressure
rfi:	Refractive Index
rinpol:	Non-polar retention indices
ripol:	Polar retention indices
tb:	Normal Boiling Point Temperature
tc:	Critical Temperature
tf:	Normal melting (fusion) point
vc:	Critical Volume
zc:	Critical Compressibility