

3-Bromobenzoic acid, 2-naphthyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C17H11BrO2/c18-15-7-3-6-14(10-15)17(19)20-16-9-8-12-4-1-2-5-13(12)11-16 |
| InchiKey: | OUYXGAYAHOBKF-UHFFFAOYSA-N |
| Formula: | C17H11BrO2 |
| SMILES: | O=C(Oc1ccc2ccccc2c1)c1cccc(Br)c1 |
| Mol. weight [g/mol]: | 327.17 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 184.87 | kJ/mol | Joback Method |
| hf | 28.51 | kJ/mol | Joback Method |
| hfus | 32.18 | kJ/mol | Joback Method |
| hvap | 76.54 | kJ/mol | Joback Method |
| log10ws | -6.53 | | Crippen Method |
| logp | 4.822 | | Crippen Method |
| mcvol | 208.350 | ml/mol | McGowan Method |
| pc | 2896.73 | kPa | Joback Method |
| rinpola | 2542.00 | | NIST Webbook |
| tb | 813.11 | K | Joback Method |
| tc | 1079.90 | K | Joback Method |
| tf | 523.89 | K | Joback Method |
| vc | 0.779 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 533.76 | J/molxK | 813.11 | Joback Method |
| cpg | 585.95 | J/molxK | 1035.44 | Joback Method |
| cpg | 577.22 | J/molxK | 990.97 | Joback Method |
| cpg | 567.77 | J/molxK | 946.51 | Joback Method |
| cpg | 557.45 | J/molxK | 902.04 | Joback Method |
| cpg | 546.16 | J/molxK | 857.58 | Joback Method |
| cpg | 594.09 | J/molxK | 1079.90 | Joback Method |
| dvisc | 0.0001834 | Paxs | 813.11 | Joback Method |
| dvisc | 0.0002180 | Paxs | 764.91 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0002653 | Paxs | 716.70 | Joback Method |
| dvisc | 0.0003321 | Paxs | 668.50 | Joback Method |
| dvisc | 0.0004304 | Paxs | 620.30 | Joback Method |
| dvisc | 0.0005827 | Paxs | 572.09 | Joback Method |
| dvisc | 0.0008342 | Paxs | 523.89 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U307555&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/17-505-2/3-Bromobenzoic-acid-2-naphthyl-ester.pdf>

Generated by Cheméo on 2024-04-19 18:26:19.609852656 +0000 UTC m=+15840428.530429978.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.