

2-Ketoglutaric acid ditbdms

Other names:	2-Ketopentanrdioic acid ditbdms 2-Ketoglutaric acid, 2tbdms derivative
Inchi:	InChI=1S/C17H34O5Si2/c1-16(2,3)23(7,8)21-14(19)12-11-13(18)15(20)22-24(9,10)17(4,
InchiKey:	RJMSHEROUAXRFK-UHFFFAOYSA-N
Formula:	C17H34O5Si2
SMILES:	CC(C)(C)[Si](C)(C)OC(=O)CCC(=O)C(=O)O[Si](C)(C)C(C)(C)C
Mol. weight [g/mol]:	374.62

Physical Properties

Property code	Value	Unit	Source
log10ws	-0.05		Crippen Method
logp	4.433		Crippen Method
rinpola	2149.00		NIST Webbook
rinpola	2149.00		NIST Webbook

Sources

Crippen Method:	http://pubs.acs.org/doi/abs/10.1021/ci9903071
Crippen Method:	https://www.chemeo.com/doc/models/crippen_log10ws
NIST Webbook:	http://webbook.nist.gov/cgi/cbook.cgi?ID=U332344&Units=SI

Legend

log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
rinpola:	Non-polar retention indices

Latest version available from:

<https://www.chemeo.com/cid/17-650-1/2-Ketoglutaric-acid-ditbdms.pdf>

Generated by Cheméo on 2024-04-28 10:10:18.263805566 +0000 UTC m=+16588267.184382894.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.