

Thymine

| | |
|----------------------|--|
| Other names: | 1-methyl-2,4(1H,3H)-pyrimidinedione 1-methyluracil 2,4(1H,3H)-Pyrimidinedione, 5-methyl- 2,4(1H,3H)-pyrimidinedione, 1-methyl- 2,4-Dihydroxy-5-methylpyrimidine 5-Methyl-2,4(1H,3H)-pyrimidinedione 5-Methyl-2,4-dioxypyrimidine 5-Methyluracil 5-methyl-2,4-dihydroxypyrimidine 5-methylpyrimidine-2,4(1H,3H)-dione N1-methyluracil Thymin |
| Inchi: | InChI=1S/C5H6N2O2/c1-3-2-6-5(9)7-4(3)8/h2H,1H3,(H2,6,7,8,9) |
| InchiKey: | RWQNBRDOKXIBIV-UHFFFAOYSA-N |
| Formula: | C5H6N2O2 |
| SMILES: | Cc1c[nH]c(=O)[nH]c1=O |
| Mol. weight [g/mol]: | 126.11 |
| CAS: | 65-71-4 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|-----------------|--------|--------------|
| affp | 880.90 | kJ/mol | NIST Webbook |
| basg | 850.00 | kJ/mol | NIST Webbook |
| chl | -2362.23 ± 0.84 | kJ/mol | NIST Webbook |
| chs | -2367.30 | kJ/mol | NIST Webbook |
| chs | -2369.00 | kJ/mol | NIST Webbook |
| ea | 0.07 ± 0.01 | eV | NIST Webbook |
| ea | 0.07 ± 0.02 | eV | NIST Webbook |
| ea | 0.06 ± 0.01 | eV | NIST Webbook |
| ea | 0.07 | eV | NIST Webbook |
| ea | 2.40 ± 0.10 | eV | NIST Webbook |
| hf | -328.70 ± 4.30 | kJ/mol | NIST Webbook |
| hfl | -462.80 ± 0.84 | kJ/mol | NIST Webbook |
| hsub | 138.00 ± 10.00 | kJ/mol | NIST Webbook |
| hsub | 131.30 ± 4.00 | kJ/mol | NIST Webbook |
| hsub | 134.10 ± 4.20 | kJ/mol | NIST Webbook |
| hvap | 134.10 ± 4.20 | kJ/mol | NIST Webbook |

| | | | |
|---------|---------------|--------|--------------------------------------|
| ie | 9.14 ± 0.03 | eV | NIST Webbook |
| ie | 9.02 | eV | NIST Webbook |
| ie | 9.20 | eV | NIST Webbook |
| ie | 9.40 ± 0.10 | eV | NIST Webbook |
| ie | 9.00 ± 0.10 | eV | NIST Webbook |
| log10ws | -1.55 | | Aqueous Solubility Prediction Method |
| log10ws | -1.51 | | Estimated Solubility Method |
| logp | -1.592 | | Crippen Method |
| mcvol | 89.250 | ml/mol | McGowan Method |
| tf | 321.30 ± 1.00 | K | NIST Webbook |
| tf | 589.53 | K | Aqueous Solubility Prediction Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|--|
| cps | 156.90 | J/molxK | 298.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 167.20 | J/molxK | 303.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 179.60 | J/molxK | 308.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |

| | | | | |
|-----|--------|---------|--------|--|
| cps | 190.90 | J/molxK | 313.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 200.50 | J/molxK | 318.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 211.10 | J/molxK | 323.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 223.00 | J/molxK | 328.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 235.00 | J/molxK | 333.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 245.90 | J/molxK | 338.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |
| cps | 256.10 | J/molxK | 343.15 | Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry |

| | | | | |
|-------|---------------|---------|--------|--|
| cps | 150.20 | J/mol×K | 298.00 | NIST Webbook |
| cps | 151.40 | J/mol×K | 298.15 | NIST Webbook |
| hfust | 17.51 | kJ/mol | 321.30 | NIST Webbook |
| hfust | 17.51 | kJ/mol | 321.30 | NIST Webbook |
| hfust | 17.51 | kJ/mol | 321.30 | NIST Webbook |
| hsubt | 125.70 ± 3.60 | kJ/mol | 410.50 | NIST Webbook |
| hsubt | 124.40 ± 1.30 | kJ/mol | 403.00 | NIST Webbook |
| psub | 3.97e-03 | kPa | 450.80 | Thermochemistry of uracil and thymine revisited |
| psub | 8.60e-04 | kPa | 430.50 | Thermochemistry of uracil and thymine revisited |
| psub | 1.34e-03 | kPa | 436.20 | Thermochemistry of uracil and thymine revisited |
| psub | 1.83e-03 | kPa | 440.50 | Thermochemistry of uracil and thymine revisited |
| psub | 3.14e-03 | kPa | 447.60 | Thermochemistry of uracil and thymine revisited |
| psub | 6.10e-04 | kPa | 426.40 | Thermochemistry of uracil and thymine revisited |
| psub | 6.01e-03 | kPa | 456.50 | Thermochemistry of uracil and thymine revisited |
| psub | 8.13e-03 | kPa | 460.60 | Thermochemistry of uracil and thymine revisited |
| psub | 0.01 | kPa | 464.80 | Thermochemistry of uracil and thymine revisited |
| psub | 0.01 | kPa | 468.90 | Thermochemistry of uracil and thymine revisited |
| psub | 0.02 | kPa | 472.30 | Thermochemistry of uracil and thymine revisited |
| rhos | 1450.00 | kg/m3 | 298.15 | Saturation molalities and standard molar enthalpies of solution of cytidine(cr), hypoxanthine(cr), thymidine(cr), thymine(cr), uridine(cr), and xanthine(cr) in H2O(l) |

Sources

Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Cyclically Polymerized Derivatives by

Differential Calorimetry: Solvation behavior of some nucleic acid bases and nucleosides in water and in aqueous guanidine

hydrochloride solutions: Viscometric, calorimetric and spectroscopic molar enthalpies of solution of cytosine(cr), hypoxanthine(cr), thymidine(cr), thymine(cr), uridine(cr), and xanthine(cr) in 90% volumetric studies on nucleic acid bases and nucleosides in aqueous guanidine hydrochloride solutions

Estimated Solubility Method:

Thermochemical study of 5-methyluracil, 6-methyluracil, and 5-nitrouracil:

<https://www.doi.org/10.1021/je060257y>

<http://onschallenge.wikispaces.com/file/view/AqueousDataset002.xlsx/351826032/AqueousDataset002.xlsx>

<https://www.doi.org/10.1016/j.jct.2015.11.029>

<http://webbook.nist.gov/cgi/cbook.cgi?ID=C65714&Units=SI>

<https://www.doi.org/10.1016/j.jct.2004.04.005>

<http://pubs.acs.org/doi/abs/10.1021/ci990307l>

<https://www.doi.org/10.1016/j.jct.2014.10.015>

<https://www.doi.org/10.1016/j.jct.2015.03.015>

<http://link.springer.com/article/10.1007/BF02311772>

http://pubs.acs.org/doi/suppl/10.1021/ci034243x/suppl_file/ci034243xsi20040112_053635.txt

<https://www.doi.org/10.1016/j.jct.2011.06.023>

Legend

| | |
|-----------------|---|
| affp: | Proton affinity |
| basg: | Gas basicity |
| chl: | Standard liquid enthalpy of combustion |
| chs: | Standard solid enthalpy of combustion |
| cps: | Solid phase heat capacity |
| ea: | Electron affinity |
| hf: | Enthalpy of formation at standard conditions |
| hfl: | Liquid phase enthalpy of formation at standard conditions |
| hfust: | Enthalpy of fusion at a given temperature |
| hsub: | Enthalpy of sublimation at standard conditions |
| hsubt: | Enthalpy of sublimation at a given temperature |
| hvap: | Enthalpy of vaporization at standard conditions |
| ie: | Ionization energy |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| psub: | Sublimation pressure |
| rhos: | Solid Density |
| tf: | Normal melting (fusion) point |

Latest version available from:

<https://www.chemeo.com/cid/18-703-1/Thymine.pdf>

Generated by Cheméo on 2025-12-23 01:47:09.579938785 +0000 UTC m=+6202627.109979450.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.