

Oxalamide

Other names:	Amid kyseliny stavelove
	Diaminoglyoxal
	Formimidic acid, 1-carbamoyl-
	NH ₂ COCONH ₂
	NSC 2770
	Oxalic acid diamide
	Oxamic acid amide
	Oxamid
	Oxamimidic acid
	ethanedioic acid, diamide
	oxamide
Inchi:	InChI=1S/C2H4N2O2/c3-1(5)2(4)6/h(H2,3,5)(H2,4,6)
InchiKey:	YIKSCQDJHCMVMK-UHFFFAOYSA-N
Formula:	C2H4N2O2
SMILES:	NC(=O)C(N)=O
Mol. weight [g/mol]:	88.07

Physical Properties

Property code	Value	Unit	Source
chs	-902.60	kJ/mol	NIST Webbook
chs	-853.43	kJ/mol	NIST Webbook
chs	-843.10 ± 4.20	kJ/mol	NIST Webbook
chs	-854.30 ± 0.40	kJ/mol	NIST Webbook
hf	-387.10 ± 1.30	kJ/mol	NIST Webbook
hfs	-456.90	kJ/mol	NIST Webbook
hfs	-515.60 ± 4.20	kJ/mol	NIST Webbook
hfs	-504.40 ± 0.50	kJ/mol	NIST Webbook
hfus	50.00	kJ/mol	Odd even effect in melting properties of 12 alkane-a,x-diamides
hsub	117.30 ± 1.20	kJ/mol	NIST Webbook
hsub	112.97	kJ/mol	NIST Webbook
ie	9.80	eV	NIST Webbook
ie	9.80	eV	NIST Webbook
ie	9.41	eV	NIST Webbook
log10ws	-0.44		Cheméo Relay (1.0)
logp	-2.043		Crippen Method

mcvol	62.140	ml/mol	McGowan Method
ss	118.11	J/mol×K	NIST Webbook
tb	473.42	K	Cheméo Relay (1.0)
tf	497.64	K	Cheméo Relay (1.0)

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	127.75	J/mol×K	497.96	Joback Method
cpg	132.85	J/mol×K	535.47	Joback Method
cpg	137.65	J/mol×K	572.98	Joback Method
cpg	142.14	J/mol×K	610.49	Joback Method
cpg	146.35	J/mol×K	648.01	Joback Method
cpg	150.26	J/mol×K	685.52	Joback Method
cpg	153.90	J/mol×K	723.03	Joback Method
cps	113.89	J/mol×K	298.15	NIST Webbook
hsubt	115.80	kJ/mol	384.00	NIST Webbook
hsubt	113.00	kJ/mol	361.00	NIST Webbook
hsubt	113.00	kJ/mol	361.00	NIST Webbook

Sources

Cheméo Relay (1.0, beta):

Odd even effect in melting properties of <https://www.doi.org/10.1016/j.jct.2006.04.004>

12 alkane-a,x-diamides:

NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=C471465&Units=SI>

Joback Method: https://en.wikipedia.org/wiki/Joback_method

McGowan Method: <http://link.springer.com/article/10.1007/BF02311772>

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci990307l>

Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws

Cheméo Relay (1.0):

Legend

chs: Standard solid enthalpy of combustion

cpg: Ideal gas heat capacity

cps:	Solid phase heat capacity
hf:	Enthalpy of formation at standard conditions
hfs:	Solid phase enthalpy of formation at standard conditions
hfus:	Enthalpy of fusion at standard conditions
hsub:	Enthalpy of sublimation at standard conditions
hsubt:	Enthalpy of sublimation at a given temperature
ie:	Ionization energy
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
ss:	Solid phase molar entropy at standard conditions
tb:	Normal Boiling Point Temperature
tf:	Normal melting (fusion) point

Latest version available from:

<https://www.cheméo.com/cid/20-251-0/Oxalamide.pdf>

Generated by Cheméo on 2026-07-21 22:33:31.262502545 +0000 UTC m=+184307.104387903.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.