

17«beta»(H),21«beta»(H)-Bishomohopanoic acid methyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C33H56O2/c1-22(10-13-28(34)35-8)23-14-19-30(4)24(23)15-20-32(6)26(30)11 |
| InchiKey: | ORWUNXLWVUEFW-PEBGAAOJSA-N |
| Formula: | C33H56O2 |
| SMILES: | COC(=O)CCC(C)C1CCC2(C)C1CCC1(C)C2CCC2C3(C)CCCC(C)(C)C3CCC21C |
| Mol. weight [g/mol]: | 484.80 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 155.77 | kJ/mol | Joback Method |
| hf | -672.99 | kJ/mol | Joback Method |
| hfus | 32.43 | kJ/mol | Joback Method |
| hvap | 91.12 | kJ/mol | Joback Method |
| log10ws | -9.31 | | Crippen Method |
| logp | 9.067 | | Crippen Method |
| mvol | 428.970 | ml/mol | McGowan Method |
| pc | 829.55 | kPa | Joback Method |
| rinpol | 3842.00 | | NIST Webbook |
| rinpol | 3842.00 | | NIST Webbook |
| tb | 1067.46 | K | Joback Method |
| tc | 1314.47 | K | Joback Method |
| tf | 685.71 | K | Joback Method |
| vc | 1.623 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|---------|---------|-----------------|---------------|
| cpg | 1803.86 | J/mol×K | 1067.46 | Joback Method |
| cpg | 1868.10 | J/mol×K | 1108.63 | Joback Method |
| cpg | 1938.24 | J/mol×K | 1149.80 | Joback Method |
| cpg | 2015.12 | J/mol×K | 1190.96 | Joback Method |
| cpg | 2099.56 | J/mol×K | 1232.13 | Joback Method |
| cpg | 2192.41 | J/mol×K | 1273.30 | Joback Method |
| cpg | 2294.48 | J/mol×K | 1314.47 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=R419019&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/21-882-9/17-beta-H-21-beta-H-Bishomohopanoic-acid-methyl-ester.pdf>

Generated by Cheméo on 2024-04-27 06:21:18.173139843 +0000 UTC m=+16488127.093717155.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.