

o-Phenanthroline

| | |
|-----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Other names: | 1,10-Fenanthroline 1,10-Phenanthroline 1,10-o-Phenanthroline 4,5-Diazaphenanthrene Activ-8 in hexylene glycol Orthophenanthroline Phenanthroline Solution forms containing 1,10-phenanthroline «beta»-Phenanthroline Â«betaÂ»-Phenanthroline |
| Inchi: | InChI=1S/C12H8N2/c1-3-9-5-6-10-4-2-8-14-12(10)11(9)13-7-1/h1-8H |
| InchiKey: | DGEZNRSVGBDHLK-UHFFFAOYSA-N |
| Formula: | C12H8N2 |
| SMILES: | c1cnc2c(c1)ccc1cccnc12 |
| Mol. weight [g/mol]: | 180.21 |
| CAS: | 66-71-7 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|-------------|--------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| basg | 908.00 | kJ/mol | NIST Webbook |
| hfus | 15.50 | kJ/mol | Thermodynamic properties of three-ring aza-aromatics. 2. Experimental results for 1,10-phenanthroline, phenanthridine, and 7,8-benzoquinoline, and mutual validation of experiments and computational methods |
| hfus | 11.80 | kJ/mol | Heat capacities and molar enthalpies and entropies of fusion for anhydrous 1,10-phenanthroline and 2,9-dimethyl-1,10-phenanthroline |
| ie | 8.51 ± 0.02 | eV | NIST Webbook |
| ie | 8.30 | eV | NIST Webbook |
| log10ws | -1.53 | | Aqueous Solubility Prediction Method |
| logp | 2.783 | | Crippen Method |
| mcvol | 137.220 | ml/mol | McGowan Method |

| | | | |
|----|--------|---|------------------------------------------------------------------------------------------------------------------|
| tf | 390.48 | K | Aqueous Solubility Prediction Method |
| tt | 390.90 | K | Thermodynamic study on six tricyclic nitrogen heterocyclic compounds by thermal analysis and effusion techniques |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------------|--------|-----------------|--------------------------------------------------------------------------------------------------|
| hfust | 15.50 | kJ/mol | 391.70 | NIST Webbook |
| hfust | 11.80 | kJ/mol | 391.10 | NIST Webbook |
| hvapt | 86.10 | kJ/mol | 298.00 | Study of the Anomalous Thermochemical Behavior of 1,2-Diazines by Correlation-Gas Chromatography |
| hvapt | 77.70 ± 0.10 | kJ/mol | 520.00 | NIST Webbook |
| hvapt | 74.90 ± 0.20 | kJ/mol | 560.00 | NIST Webbook |
| hvapt | 72.10 ± 0.20 | kJ/mol | 600.00 | NIST Webbook |

Sources

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=C66717&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Thermodynamic properties of three-ring aza-aromatics. 2. Experimental results for molar enthalpies and entropies of fusion for anhydrous 1,2-diazines and their corresponding neutral and cationic forms. | https://www.doi.org/10.1016/j.jct.2009.11.011 |
| Experimental results for molar enthalpies and entropies of fusion for anhydrous 1,2-diazines and their corresponding neutral and cationic forms. | https://www.doi.org/10.1016/j.tca.2007.10.001 |
| Thermodynamic study on six tricyclic nitrogen heterocyclic compounds by thermal analysis and effusion techniques. | https://www.doi.org/10.1016/j.tca.2016.05.001 |
| Thermal analysis and effusion techniques. | https://www.doi.org/10.1021/je900702t |
| Aqueous Solubility Prediction Method: Chromatography. | http://onschallenge.wikispaces.com/file/view/AqueousDataset002.xlsx/351826032/AqueousDataset002.xlsx |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |

Legend

| | |
|--------------|-------------------------------------------|
| basg: | Gas basicity |
| hfus: | Enthalpy of fusion at standard conditions |

| | |
|-----------------|-------------------------------------------------|
| hfust: | Enthalpy of fusion at a given temperature |
| hvapt: | Enthalpy of vaporization at a given temperature |
| ie: | Ionization energy |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| tf: | Normal melting (fusion) point |
| tt: | Triple Point Temperature |

Latest version available from:

<https://www.chemeo.com/cid/27-145-1/o-Phenanthroline.pdf>

Generated by Cheméo on 2024-04-25 08:03:15.397549011 +0000 UTC m=+16321444.318126323.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.