

«beta»-N-Acetyl-D-glucosamine, TMS

Inchi: InChI=1S/C19H45NO6Si4/c1-14(21)20-15-16(23-27(2,3)4)17(24-28(5,6)7)19(26-30(11,12)13)22-18-25-29(8,9)10-3
InchiKey: HUXJBHUJRMWUPT-QTMHVTGLSA-N
Formula: C19H45NO6Si4
SMILES: CC(=O)NC1C(O[Si](C)(C)C)OC(O[Si](C)(C)C)C(O[Si](C)(C)C)C1O[Si](C)(C)C
Mol. weight [g/mol]: 495.91

Physical Properties

Property code	Value	Unit	Source
log10ws	4.18		Crippen Method
logp	4.317		Crippen Method
rinpol	2144.00		NIST Webbook

Sources

Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws
NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=R441100&Units=SI>
Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci990307l>

Legend

log10ws: Log10 of Water solubility in mol/l
logp: Octanol/Water partition coefficient
rinpol: Non-polar retention indices

Latest version available from:

<https://www.chemeo.com/cid/27-709-5/beta-N-Acetyl-D-glucosamine-TMS.pdf>

Generated by Cheméo on 2025-12-05 13:54:48.235238358 +0000 UTC m=+4691085.765279013.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.