

«beta»-Ionone, 5,6-epoxy

Inchi:	InChI=1S/C13H20O2/c1-10(14)6-9-13-11(2,3)7-5-8-12(13,4)15-13/h6,9H,5,7-8H2,1-4H3
InchiKey:	ZTJZJYUGOJYHCU-RMKNXTFCSA-N
Formula:	C13H20O2
SMILES:	CC(=O)C=CC1C(C)(C)CCC2OC21C
Mol. weight [g/mol]:	208.30

Physical Properties

Property code	Value	Unit	Source
hfus	22.92	kJ/mol	Joback Method
logp	2.725		Crippen Method
mcvol	175.450	ml/mol	McGowan Method
rinpol	1488.00		NIST Webbook

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	474.18	J/mol×K	590.71	Joback Method
cpg	492.47	J/mol×K	627.68	Joback Method
cpg	509.60	J/mol×K	664.66	Joback Method
cpg	525.85	J/mol×K	701.63	Joback Method
cpg	541.47	J/mol×K	738.60	Joback Method
cpg	556.73	J/mol×K	775.57	Joback Method
cpg	571.91	J/mol×K	812.55	Joback Method

Sources

Joback Method:	https://en.wikipedia.org/wiki/Joback_method
McGowan Method:	http://link.springer.com/article/10.1007/BF02311772
Crippen Method:	http://pubs.acs.org/doi/abs/10.1021/ci990307l
Crippen Method:	https://www.cheméo.com/doc/models/crippen_log10ws
Cheméo Relay (1.0):	

Cheméo Relay (1.0, beta):

NIST Webbook:

<http://webbook.nist.gov/cgi/cbook.cgi?ID=R568839&Units=SI>

Legend

cpg:	Ideal gas heat capacity
hfus:	Enthalpy of fusion at standard conditions
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
rinpol:	Non-polar retention indices

Latest version available from:

<https://www.chemeo.com/cid/28-102-7/beta-Ionone-5-6-epoxy.pdf>

Generated by Cheméo on 2026-07-22 01:21:18.535189447 +0000 UTC m=+194374.377074827.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.