

(p-methylbenzylidene)-octyl-amine

Inchi: InChI=1S/C16H25N/c1-3-4-5-6-7-8-13-17-14-16-11-9-15(2)10-12-16/h9-12,14H,3-8,13H2
InchiKey: DUYWJSXROTZXSC-SAPNQHFASA-N
Formula: C16H25N
SMILES: CCCCCCCN=Cc1ccc(C)cc1
Mol. weight [g/mol]: 231.38

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|--------|----------------|
| hf | -66.29 | kJ/mol | Joback Method |
| hvap | 57.46 | kJ/mol | Joback Method |
| log10ws | -4.95 | | Crippen Method |
| logp | 4.774 | | Crippen Method |
| mcvol | 218.220 | ml/mol | McGowan Method |
| pc | 1546.35 | kPa | Joback Method |
| rinpol | 1881.00 | | NIST Webbook |
| tb | 673.82 | K | Joback Method |
| tc | 878.02 | K | Joback Method |

Sources

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>
Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws
Joback Method: https://en.wikipedia.org/wiki/Joback_method
McGowan Method: <http://link.springer.com/article/10.1007/BF02311772>
NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=R160391&Units=SI>

Legend

hf: Enthalpy of formation at standard conditions
hvap: Enthalpy of vaporization at standard conditions
log10ws: Log10 of Water solubility in mol/l

| | |
|----------------|-------------------------------------|
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |

Latest version available from:

<https://www.cheméo.com/cid/37-003-7/p-methylbenzylidene-octyl-amine.pdf>

Generated by Cheméo on 2024-04-19 19:52:40.507150347 +0000 UTC m=+15845609.427727662.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.