

Hexanoic acid, 6-(carboxyamino)-, n-benzyl ester

| | |
|-----------------------------|---|
| Other names: | 6-[[[(phenylmethoxy)carbonyl]amino]hexanoic acid |
| Inchi: | InChI=1S/C14H19NO4/c16-13(17)9-5-2-6-10-15-14(18)19-11-12-7-3-1-4-8-12/h1,3-4,7-8 |
| InchiKey: | RXQDBVWDABAAHL-UHFFFAOYSA-N |
| Formula: | C14H19NO4 |
| SMILES: | O=C(O)CCCCNC(=O)OCc1ccccc1 |
| Mol. weight [g/mol]: | 265.31 |
| CAS: | 1947-00-8 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -230.86 | kJ/mol | Joback Method |
| hf | -551.90 | kJ/mol | Joback Method |
| hfus | 39.63 | kJ/mol | Joback Method |
| hvap | 88.05 | kJ/mol | Joback Method |
| log10ws | -3.41 | | Crippen Method |
| logp | 2.558 | | Crippen Method |
| mcvol | 209.220 | ml/mol | McGowan Method |
| pc | 2458.04 | kPa | Joback Method |
| tb | 818.91 | K | Joback Method |
| tc | 1019.78 | K | Joback Method |
| tf | 509.53 | K | Joback Method |
| vc | 0.795 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 622.14 | J/molxK | 818.91 | Joback Method |
| cpg | 633.71 | J/molxK | 852.39 | Joback Method |
| cpg | 644.45 | J/molxK | 885.87 | Joback Method |
| cpg | 654.38 | J/molxK | 919.35 | Joback Method |
| cpg | 663.53 | J/molxK | 952.83 | Joback Method |
| cpg | 671.94 | J/molxK | 986.30 | Joback Method |
| cpg | 679.64 | J/molxK | 1019.78 | Joback Method |

Sources

| | |
|------------------------|---|
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=C1947008&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| h vap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/45-411-5/Hexanoic-acid-6-carboxyamino-n-benzyl-ester.pdf>

Generated by Cheméo on 2024-04-24 09:36:04.376145935 +0000 UTC m=+16240613.296723256.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.