Benzoic acid, 2-iodo-

Other names:	2-Iodobenzoic acid
	Benzoic acid, o-iodo-
	Kyselina o-jodbenzoova
	USAF ek-572
	o-Iodobenzoic acid
Inchi:	InChI=1S/C7H5IO2/c8-6-4-2-1-3-5(6)7(9)10/h1-4H,(H,9,10)
InchiKey:	CJNZAXGUTKBIHP-UHFFFAOYSA-N
Formula:	C7H5IO2
SMILES:	O=C(O)c1ccccc1I
Mol. weight [g/mol]:	248.02
CAS:	88-67-5

Physical Properties

Property code	Value	Unit	Source
chs	-3173.70 ± 1.70	kJ/mol	NIST Webbook
chs	-3228.80 ± 7.20	kJ/mol	NIST Webbook
chs	-3167.00 ± 2.00	kJ/mol	NIST Webbook
gf	-96.78	kJ/mol	Joback Method
hf	-202.80 ± 1.70	kJ/mol	NIST Webbook
hfs	-295.40 ± 1.70	kJ/mol	NIST Webbook
hfus	17.63	kJ/mol	Joback Method
hsub	103.00 ± 0.40	kJ/mol	NIST Webbook
hsub	112.80 ± 2.00	kJ/mol	NIST Webbook
hsub	92.60 ± 0.20	kJ/mol	NIST Webbook
hvap	66.91	kJ/mol	Joback Method
log10ws	-2.73		Aqueous Solubility Prediction Method
logp	1.989		Crippen Method
mcvol	118.990	ml/mol	McGowan Method
pc	4815.84	kPa	Joback Method
tb	630.41	K	Joback Method
tc	868.95	K	Joback Method
tf	434.50	K	Thermochemistry of halogenobenzoic acids as an access to PC-SAFT solubility modeling
tf	435.53	K	Aqueous Solubility Prediction Method
tt	435.02 ± 0.02	K	NIST Webbook

vc	0.432	m3/kmol	Joback Method

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	262.84	J/mol×K	868.95	Joback Method
cpg	227.76	J/mol×K	630.41	Joback Method
cpg	234.94	J/mol×K	670.17	Joback Method
cpg	241.53	J/mol×K	709.92	Joback Method
cpg	247.57	J/mol×K	749.68	Joback Method
cpg	253.10	J/mol×K	789.44	Joback Method
cpg	258.18	J/mol×K	829.20	Joback Method
dvisc	0.0001104	Pa×s	630.41	Joback Method
dvisc	0.0036678	Pa×s	376.40	Joback Method
dvisc	0.0015227	Pa×s	418.74	Joback Method
dvisc	0.0007429	Pa×s	461.07	Joback Method
dvisc	0.0004089	Pa×s	503.41	Joback Method
dvisc	0.0002470	Pa×s	545.74	Joback Method
dvisc	0.0001604	Pa×s	588.08	Joback Method
hfust	21.38	kJ/mol	435.10	NIST Webbook
hfust	21.38	kJ/mol	435.10	NIST Webbook
hsubt	111.40 ± 0.80	kJ/mol	352.00	NIST Webbook

Sources

Aqueous Solubility Prediction Method:	http://onschallenge.wikispaces.com/file/view/AqueousDataset002.xlsx/351826032/AqueousDa
McGowan Method:	http://link.springer.com/article/10.1007/BF02311772
NIST Webbook:	http://webbook.nist.gov/cgi/cbook.cgi?ID=C88675&Units=SI
Crippen Method:	http://pubs.acs.org/doi/abs/10.1021/ci990307l
Thermochemistry of halogenobenzoic acids as an access to PC-SAFT solubility modeling:	https://www.doi.org/10.1016/j.fluid.2015.10.001
Joback Method:	https://en.wikipedia.org/wiki/Joback_method

Legend

chs: Standard solid enthalpy of combustion

cpg:	Ideal gas heat capacity
dvisc:	Dynamic viscosity
gf:	Standard Gibbs free energy of formation
hf:	Enthalpy of formation at standard conditions
hfs:	Solid phase enthalpy of formation at standard conditions
hfus:	Enthalpy of fusion at standard conditions
hfust:	Enthalpy of fusion at a given temperature
hsub:	Enthalpy of sublimation at standard conditions
hsubt:	Enthalpy of sublimation at a given temperature
hvap:	Enthalpy of vaporization at standard conditions
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
pc:	Critical Pressure
tb:	Normal Boiling Point Temperature
tc:	Critical Temperature
tf:	Normal melting (fusion) point
tt:	Triple Point Temperature
vc:	Critical Volume

Latest version available from:

https://www.chemeo.com/cid/47-162-0/Benzoic-acid-2-iodo.pdf

Generated by Cheméo on 2024-10-12 05:05:55.931621324 +0000 UTC m=+3290418.568590622.

Cheméo (https://www.chemeo.com) is the biggest free database of chemical and physical data for the process industry.