

Italicen-12-yl acetate

| | |
|-----------------------------|---|
| Inchi: | InChI=1S/C17H26O2/c1-11-7-8-17-12(2)5-6-14(17)16(4,15(17)9-11)10-19-13(3)18/h9,12 |
| InchiKey: | WEGYMQPVQUNDTQ-VFTHGPRRSA-N |
| Formula: | C17H26O2 |
| SMILES: | CC(=O)OCC1(C)C2C=C(C)CCC23C(C)CCC13 |
| Mol. weight [g/mol]: | 262.39 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 10.32 | kJ/mol | Joback Method |
| hf | -396.82 | kJ/mol | Joback Method |
| hfus | 23.16 | kJ/mol | Joback Method |
| hvap | 60.71 | kJ/mol | Joback Method |
| log10ws | -4.13 | | Crippen Method |
| logp | 3.958 | | Crippen Method |
| mcvol | 220.950 | ml/mol | McGowan Method |
| pc | 1859.51 | kPa | Joback Method |
| rinsol | 1780.00 | | NIST Webbook |
| tb | 688.69 | K | Joback Method |
| tc | 909.97 | K | Joback Method |
| tf | 452.89 | K | Joback Method |
| vc | 0.847 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 668.83 | J/mol×K | 688.69 | Joback Method |
| cpg | 689.94 | J/mol×K | 725.57 | Joback Method |
| cpg | 710.27 | J/mol×K | 762.45 | Joback Method |
| cpg | 730.07 | J/mol×K | 799.33 | Joback Method |
| cpg | 749.63 | J/mol×K | 836.21 | Joback Method |
| cpg | 769.19 | J/mol×K | 873.09 | Joback Method |
| cpg | 789.03 | J/mol×K | 909.97 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=R233162&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvac: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mccol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/47-399-8/Italicen-12-yl-acetate.pdf>

Generated by Cheméo on 2024-04-25 15:33:39.186590603 +0000 UTC m=+16348468.107167918.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.