

2-Butenoic acid, 2-methyl-, 2,3,5,7a-tetrahydro-7-(hydroxymethyl)-1H-pyrrolizine ester, [1S-[1«alpha»(Z),7a«alpha»]]-

Other names:

Heliotridine, 7-(2-methylbutanoate), (Z)
Heliotridine, 7-angelate (ester)

Rivularine

Heliotridine, 7-angelyl-

7-Angeloylheliotridine

7-Angelylheliotridine

7-(Hydroxymethyl)-(1S,7aR)-2,3,5,7a-tetrahydro-1H-pyrrolizin-1-yl
(2Z)-2-methyl-2-butenoate
Heliotridine, 7-angelate

Inchi: InChI=1S/C13H19NO3/c1-3-9(2)13(16)17-11-5-7-14-6-4-10(8-15)12(11)14/h3-4,11-12,15
InchiKey: TYGYPIIOOQNWBU-YKSZOPSQSA-N
Formula: C13H19NO3
SMILES: CC=C(C)C(=O)OC1CCN2CC=C(CO)C12
Mol. weight [g/mol]: 237.29
CAS: 723-78-4

Physical Properties

Property code	Value	Unit	Source
log10ws	-1.68		Crippen Method
logp	0.871		Crippen Method
mcvol	187.000	ml/mol	McGowan Method
rinpol	1820.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1818.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1818.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1820.00		NIST Webbook
rinpol	1821.00		NIST Webbook

Sources

McGowan Method:

<http://link.springer.com/article/10.1007/BF02311772>

NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=C723784&Units=SI>
Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>
Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws

Legend

log10ws: Log10 of Water solubility in mol/l
logp: Octanol/Water partition coefficient
mcvol: McGowan's characteristic volume
rinpolar: Non-polar retention indices

Latest version available from:

<https://www.chemeo.com/cid/47-524-8/2-Butenoic-acid-2-methyl-2-3-5-7a-tetrahydro-7-hydroxymethyl-1H-pyrrolizin-1>

Generated by Cheméo on 2024-04-25 19:03:22.170342898 +0000 UTC m=+16361051.090920213.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.