

1,8-Nonadiene, 2,4,6,8-tetramethyl, # 1

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C13H24/c1-10(2)7-12(5)9-13(6)8-11(3)4/h12-13H,1,3,7-9H2,2,4-6H3 |
| InchiKey: | ZWOKFAVFJDZGAW-UHFFFAOYSA-N |
| Formula: | C13H24 |
| SMILES: | C=C(C)CC(C)CC(C)CC(=C)C |
| Mol. weight [g/mol]: | 180.33 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 212.28 | kJ/mol | Joback Method |
| hf | -90.93 | kJ/mol | Joback Method |
| hfus | 17.20 | kJ/mol | Joback Method |
| hvap | 42.58 | kJ/mol | Joback Method |
| log10ws | -4.49 | | Crippen Method |
| logp | 4.581 | | Crippen Method |
| mcvol | 185.430 | ml/mol | McGowan Method |
| pc | 1801.56 | kPa | Joback Method |
| rinpol | 1154.00 | | NIST Webbook |
| rinpol | 1154.00 | | NIST Webbook |
| tb | 489.08 | K | Joback Method |
| tc | 667.82 | K | Joback Method |
| tf | 174.83 | K | Joback Method |
| vc | 0.716 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 418.47 | J/mol×K | 489.08 | Joback Method |
| cpg | 436.41 | J/mol×K | 518.87 | Joback Method |
| cpg | 453.54 | J/mol×K | 548.66 | Joback Method |
| cpg | 469.88 | J/mol×K | 578.45 | Joback Method |
| cpg | 485.47 | J/mol×K | 608.24 | Joback Method |
| cpg | 500.33 | J/mol×K | 638.03 | Joback Method |
| cpg | 514.49 | J/mol×K | 667.82 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=R568111&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307I |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinp: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/48-996-4/1-8-Nonadiene-2-4-6-8-tetramethyl-1.pdf>

Generated by Cheméo on 2024-04-27 07:42:17.531524941 +0000 UTC m=+16492986.452102253.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.