

Fumaric acid, monoamide, N,N-dimethyl-, 2-naphthyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C16H15NO3/c1-17(2)15(18)9-10-16(19)20-14-8-7-12-5-3-4-6-13(12)11-14/h3- |
| InchiKey: | BYLYCUBVNBFWBF-MDZDMLPSA-N |
| Formula: | C16H15NO3 |
| SMILES: | CN(C)C(=O)C=CC(=O)Oc1ccc2ccccc2c1 |
| Mol. weight [g/mol]: | 269.30 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 121.43 | kJ/mol | Joback Method |
| hf | -130.07 | kJ/mol | Joback Method |
| hfus | 35.48 | kJ/mol | Joback Method |
| hvap | 73.69 | kJ/mol | Joback Method |
| log10ws | -3.46 | | Crippen Method |
| logp | 2.390 | | Crippen Method |
| mcvol | 207.770 | ml/mol | McGowan Method |
| pc | 2417.12 | kPa | Joback Method |
| rinpol | 2679.00 | | NIST Webbook |
| rinpol | 2679.00 | | NIST Webbook |
| tb | 762.88 | K | Joback Method |
| tc | 991.96 | K | Joback Method |
| tf | 491.20 | K | Joback Method |
| vc | 0.773 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 567.36 | J/mol×K | 762.88 | Joback Method |
| cpg | 580.73 | J/mol×K | 801.06 | Joback Method |
| cpg | 593.10 | J/mol×K | 839.24 | Joback Method |
| cpg | 604.58 | J/mol×K | 877.42 | Joback Method |
| cpg | 615.25 | J/mol×K | 915.60 | Joback Method |
| cpg | 625.20 | J/mol×K | 953.78 | Joback Method |
| cpg | 634.53 | J/mol×K | 991.96 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U357439&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinp: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/49-588-6/Fumaric-acid-monoamide-N-N-dimethyl-2-naphthyl-ester.pdf>

Generated by Cheméo on 2024-05-02 11:17:03.53838782 +0000 UTC m=+16937872.458965135.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.