

Glycine, n-(2-furanylcarbonyl)-, methyl ester

Other names:	Glycine, N-2-furoyl-, methyl ester Methyl N-2-furoylglycinate Methyl (2-furoylamino)acetate N-(2-Furoyl)glycine, methyl ester [(Furan-2-carbonyl)-amino]-acetic acid methyl ester
Inchi:	InChI=1S/C8H9NO4/c1-12-7(10)5-9-8(11)6-3-2-4-13-6/h2-4H,5H2,1H3,(H,9,11)
InchiKey:	MPRFBNLQEYNYQK-UHFFFAOYSA-N
Formula:	C8H9NO4
SMILES:	COC(=O)CNC(=O)c1ccco1
Mol. weight [g/mol]:	183.16

Physical Properties

Property code	Value	Unit	Source
hf	-540.86	kJ/mol	Cheméo Relay (1.0)
hvap	75.22	kJ/mol	Cheméo Relay (1.0)
ie	8.95	eV	Cheméo Relay (1.0)
log10ws	-1.51		Cheméo Relay (1.0)
logp	0.182		Crippen Method
mcvol	128.980	ml/mol	McGowan Method
rinpol	1523.60		NIST Webbook
rinpol	1471.00		NIST Webbook
rinpol	1471.00		NIST Webbook
rinpol	1471.00		NIST Webbook
rinpol	1522.20		NIST Webbook
tb	561.65	K	Cheméo Relay (1.0)
tf	367.99	K	Cheméo Relay (1.0)

Sources

Cheméo Relay (1.0, beta):

NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=C13290001&Units=SI>

McGowan Method: <http://link.springer.com/article/10.1007/BF02311772>

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>

Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws

Legend

hf:	Enthalpy of formation at standard conditions
hvap:	Enthalpy of vaporization at standard conditions
ie:	Ionization energy
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
rlnpol:	Non-polar retention indices
tb:	Normal Boiling Point Temperature
tf:	Normal melting (fusion) point

Latest version available from:

<https://www.chemeo.com/cid/54-671-7/Glycine-n-2-furanylcarbonyl-methyl-ester.pdf>

Generated by Cheméo on 2026-07-22 02:14:34.325838604 +0000 UTC m=+197570.167723996.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.