5,6-Dihydro-6-methyluracil

Other names: 2,4(1H,3H)-Pyrimidinedione, dihydro-6-methyl-

4-Methyldihydrouracil

5,6-dihydro-5-methyluracil

5,6-dihydrothymine

5-methyl-5,6-dihydrouracil

5-methyldihydropyrimidine-2,4(1H,3H)-dione

5-methylhydrouracil

6-Methyl-5,6-dihydrouracil

6-Methyldihydro-2,4(1H,3H)-pyrimidinedione

6-Methyldihydrouracil Dihydro-6-methyluracil Hydrouracil, 6-methyl-

Metacil, dihydro-

NSC 44127

InChl=1S/C5H8N2O2/c1-3-2-4(8)7-5(9)6-3/h3H,2H2,1H3,(H2,6,7,8,9)

InchiKey: XQLIRTZXJDEQAO-UHFFFAOYSA-N

Formula: C5H8N2O2

SMILES: CC1CC(=O)NC(=O)N1

Mol. weight [g/mol]: 128.13 CAS: 2434-49-3

Physical Properties

| Property code | Value | Unit | Source | |
|---------------|----------|---------|----------------|--|
| chs | -2592.00 | kJ/mol | NIST Webbook | |
| chs | -2592.40 | kJ/mol | NIST Webbook | |
| gf | -54.09 | kJ/mol | Joback Method | |
| hf | -291.99 | kJ/mol | Joback Method | |
| hfus | 18.74 | kJ/mol | Joback Method | |
| hvap | 49.16 | kJ/mol | Joback Method | |
| log10ws | -0.83 | | Crippen Method | |
| logp | -0.396 | | Crippen Method | |
| mcvol | 93.550 | ml/mol | McGowan Method | |
| рс | 5304.68 | kPa | Joback Method | |
| tb | 566.09 | K | Joback Method | |
| tc | 826.71 | K | Joback Method | |
| tf | 499.99 | K | Joback Method | |
| VC | 0.337 | m3/kmol | Joback Method | |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [| [K] Source | |
|---------------|----------|---------|---------------|--|--|
| cpg | 295.01 | J/mol×K | 826.71 | Joback Method | |
| cpg | 284.87 | J/mol×K | 783.28 | Joback Method | |
| cpg | 273.71 | J/mol×K | 739.84 | Joback Method | |
| cpg | 261.65 | J/mol×K | 696.40 | Joback Method | |
| cpg | 248.80 | J/mol×K | 652.96 | Joback Method | |
| cpg | 235.27 | J/mol×K | 609.53 | Joback Method | |
| cpg | 221.19 | J/mol×K | 566.09 | Joback Method | |
| psub | 7.56e-04 | kPa | | Enthalpies of formation of 6,6-dihydro-5-methylurad and 6,6-dihydro-6-methylurad | |
| psub | 6.32e-04 | kPa | | Enthalpies of formation of 6,6-dihydro-5-methylurac and 6,6-dihydro-6-methylurac | |
| psub | 7.86e-04 | kPa | | Enthalpies of formation of 6,6-dihydro-5-methylurac and 6,6-dihydro-6-methylurac | |
| psub | 9.06e-04 | kPa | | Enthalpies of formation of 5,6-dihydro-5-methylurac and 5,6-dihydro-6-methylurac | |
| psub | 1.11e-03 | kPa | | Enthalpies of formation of i,6-dihydro-5-methylurac and i,6-dihydro-6-methylurac | |
| psub | 1.31e-04 | kPa | | Enthalpies of formation of i,6-dihydro-5-methylurac and i,6-dihydro-6-methylurac | |
| psub | 1.63e-04 | kPa | | Enthalpies of formation of 6,6-dihydro-5-methylurac and 6,6-dihydro-6-methylurac | |
| psub | 1.97e-04 | kPa | | Enthalpies of formation of i,6-dihydro-5-methylurad and i,6-dihydro-6-methylurad | |

| psub | 2.42e-04 | kPa | 385.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
|------|-----------------------|-----|--------|--|
| psub | 2.98e-04 | kPa | 387.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| | | | | 5,6-dihydro-6-methyluracil |
| psub | 3.59e-04 | kPa | 389.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| | | | | 5,6-dihydro-6-methyluracil |
| psub | 4.27e-04 | kPa | 391.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro 6 methyluracil |
| nguh | 5.21e-04 | kPa | 393.19 | 5,6-dihydro-6-methyluracil |
| psub | 5.21 e- 04 | кра | 393.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 6.33e-04 | kPa | 395.16 | Enthalpies of |
| рэць | 0.006-04 | NIα | 393.10 | formation of 5,6-dihydro-5-methyluracil and |
| | | | | 5,6-dihydro-6-methyluracil |
| psub | 5.38e-04 | kPa | 393.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 9.51e-04 | kPa | 399.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.10e-03 | kPa | 401.16 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.27e-04 | kPa | 379.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.57e-04 | kPa | 381.18 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.90e-04 | kPa | 383.14 | Enthalpies of formation of 5,6-dihydro-5-methyluracil |
| | | | | and 5,6-dihydro-6-methyluracil |

| | psub | 2.34e-04 | kPa | 385.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6 methyluracil |
|---|------|-----------|-----|--------|--|
| | psub | 2.87e-04 | kPa | 387.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| | | | | | 5,6-dihydro-6-methyluracil |
| | psub | 3.46e-04 | kPa | 389.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| _ | | | | | 5,6-dihydro-6-methyluracil |
| | psub | 4.20e-04 | kPa | 391.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| | nauh | 5.08e-04 | kPa | 393.19 | 5,6-dihydro-6-methyluracil |
| | psub | 5.00e-04 | кра | 393.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| | psub | 6.13e-04 | kPa | 395.16 | Enthalpies of |
| | pour | 0.100 0 1 | & | 000110 | formation of 5,6-dihydro-5-methyluracil and |
| | | | | | 5,6-dihydro-6-methyluracil |
| | psub | 7.37e-04 | kPa | 397.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| | psub | 8.82e-04 | kPa | 399.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| | psub | 1.07e-03 | kPa | 401.16 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| | psub | 4.42e-04 | kPa | 391.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| | psub | 3.64e-04 | kPa | 389.15 | Enthalpies of formation of 5,6-dihydro-5-methyluracil |
| | | | | | and 5,6-dihydro-6-methyluracil |
| | psub | 3.01e-04 | kPa | 387.19 | Enthalpies of formation of 5,6-dihydro-5-methyluracil and |
| | | | | | 5,6-dihydro-6-methyluracil |

| psub | 2.43e-04 | kPa | 385.15 Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
|------|----------|-----|---|
| psub | 1.98e-04 | kPa | 383.14 Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.68e-04 | kPa | 381.18 Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |
| psub | 1.38e-04 | kPa | 379.15 Enthalpies of formation of 5,6-dihydro-5-methyluracil and 5,6-dihydro-6-methyluracil |

Sources

Crippen Method: http://pubs.acs.org/doi/abs/10.1021/ci990307l

Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws

Enthalpies of formation of 5,6-dihydro-5-methyluracil and spantylled priethyluracil: https://www.doi.org/10.1016/j.jct.2013.05.002 https://en.wikipedia.org/wiki/Joback_method

McGowan Method: http://link.springer.com/article/10.1007/BF02311772

NIST Webbook: http://webbook.nist.gov/cgi/cbook.cgi?ID=C2434493&Units=SI

Legend

chs: Standard solid enthalpy of combustion

cpg: Ideal gas heat capacity

gf: Standard Gibbs free energy of formationhf: Enthalpy of formation at standard conditionshfus: Enthalpy of fusion at standard conditions

hvap: Enthalpy of vaporization at standard conditions

log10ws: Log10 of Water solubility in mol/llogp: Octanol/Water partition coefficientmcvol: McGowan's characteristic volume

pc: Critical Pressurepsub: Sublimation pressure

tb: Normal Boiling Point Temperature

tc: Critical Temperature

tf: Normal melting (fusion) point

vc: Critical Volume

Latest version available from:

https://www.chemeo.com/cid/56-905-5/5-6-Dihydro-6-methyluracil.pdf

Generated by Cheméo on 2024-04-20 06:20:26.227960275 +0000 UTC m=+15883275.148537587.

Cheméo (https://www.chemeo.com) is the biggest free database of chemical and physical data for the process industry.