

1-Methylthymine

Other names:	1,5-dimethyluracil
Inchi:	InChI=1S/C6H8N2O2/c1-4-3-8(2)6(10)7-5(4)9/h3H,1-2H3,(H,7,9,10)
InchiKey:	GKMIDMKPBOUSBQ-UHFFFAOYSA-N
Formula:	C6H8N2O2
SMILES:	Cc1cn(C)c(=O)[nH]c1=O
Mol. weight [g/mol]:	140.14
CAS:	4160-72-9

Physical Properties

Property code	Value	Unit	Source
ea	0.04 ± 0.01	eV	NIST Webbook
log10ws	-1.30		Crippen Method
logp	-1.100		Crippen Method
mcvol	103.340	ml/mol	McGowan Method

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cps	187.60	J/molxK	298.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	190.00	J/molxK	303.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry

cps	190.30	J/molxK	308.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	194.10	J/molxK	313.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	194.80	J/molxK	318.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	198.50	J/molxK	323.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	199.20	J/molxK	328.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	202.00	J/molxK	333.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	203.00	J/molxK	338.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry

cps	205.20	J/mol×K	343.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
hsubt	124.40 ± 1.30	kJ/mol	403.00	NIST Webbook

Sources

Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry:
NIST Webbook:

<https://www.doi.org/10.1021/je060257y>

<http://link.springer.com/article/10.1007/BF02311772>

<http://webbook.nist.gov/cgi/cbook.cgi?ID=C4160729&Units=SI>

Crippen Method:

<http://pubs.acs.org/doi/abs/10.1021/ci990307l>

Crippen Method:

https://www.chemeo.com/doc/models/crippen_log10ws

Legend

cps:	Solid phase heat capacity
ea:	Electron affinity
hsubt:	Enthalpy of sublimation at a given temperature
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume

Latest version available from:

<https://www.chemeo.com/cid/66-065-7/1-Methylthymine.pdf>

Generated by Cheméo on 2025-12-05 07:42:42.587486803 +0000 UTC m=+4668760.117527466.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.