

Neodihydrocarveol

| | |
|-----------------------------|--|
| Other names: | neo-iso-Dihydrocarveol |
| Inchi: | InChI=1S/C10H18O/c1-7(2)9-5-4-8(3)10(11)6-9/h8-11H,1,4-6H2,2-3H3 |
| InchiKey: | KRCZYMFUWVJCLI-UHFFFAOYSA-N |
| Formula: | C10H18O |
| SMILES: | <chem>C=C(C)C1CCC(C)C(O)C1</chem> |
| Mol. weight [g/mol]: | 154.25 |
| CAS: | 18675-34-8 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|---------|----------------|
| gf | -15.18 | kJ/mol | Joback Method |
| hf | -272.68 | kJ/mol | Joback Method |
| hfus | 17.13 | kJ/mol | Joback Method |
| hvap | 53.75 | kJ/mol | Joback Method |
| log10ws | -2.65 | | Crippen Method |
| logp | 2.360 | | Crippen Method |
| mcvol | 142.470 | ml/mol | McGowan Method |
| pc | 2770.08 | kPa | Joback Method |
| tb | 527.15 | K | Joback Method |
| tc | 719.80 | K | Joback Method |
| tf | 246.46 | K | Joback Method |
| vc | 0.527 | m3/kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 350.71 | J/molxK | 527.15 | Joback Method |
| cpg | 367.51 | J/molxK | 559.26 | Joback Method |
| cpg | 383.50 | J/molxK | 591.37 | Joback Method |
| cpg | 398.69 | J/molxK | 623.47 | Joback Method |
| cpg | 413.11 | J/molxK | 655.58 | Joback Method |
| cpg | 426.76 | J/molxK | 687.69 | Joback Method |
| cpg | 439.67 | J/molxK | 719.80 | Joback Method |

Sources

| | |
|------------------------|---|
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=C18675348&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| h vap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/66-602-0/Neodihydrocarveol.pdf>

Generated by Cheméo on 2024-04-28 18:14:05.42529204 +0000 UTC m=+16617294.345869362.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.