1,3-Dimethyl-5-ethyluracil

Other names: 2,4(1H,3H)-pyrimidinedione, 5-ethyl-1,3-dimethyl-

uracil, 5-ethyl-1,3-dimethyl-

Inchi: InChl=1S/C8H12N2O2/c1-4-6-5-9(2)8(12)10(3)7(6)11/h5H,4H2,1-3H3

InchiKey: RGFNQXKFHSVDHI-UHFFFAOYSA-N

Formula: C8H12N2O2

SMILES: CCc1cn(C)c(=O)n(C)c1=O

Mol. weight [g/mol]: 168.19 CAS: 31703-08-9

Physical Properties

Property code	Value	Unit	Source
log10ws	-3.69		Crippen Method
logp	-0.354		Crippen Method
mcvol	131.520	ml/mol	McGowan Method

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]] Source
cps	275.20	J/mol×K	333.15 Cy	Heat Capacities of Uracil, Thymine, and Its Alkylated, clooligomethylenated and Halogenated Derivatives by Differential Calorimetry
cps	291.10	J/mol×K	343.15 Cy	Heat Capacities of Uracil, Thymine, and Its Alkylated, clooligomethylenated and Halogenated Derivatives by Differential Calorimetry

cps	281.70	J/mol×K	338.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	241.10	J/mol×K	298.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	250.10	J/mol×K	303.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	256.50	J/mol×K	308.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	261.00	J/mol×K	313.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	264.30	J/mol×K	318.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
cps	267.40	J/mol×K	323.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry

cps	270.70	J/mol×K	328.15	Heat Capacities of Uracil, Thymine, and Its Alkylated, Cyclooligomethylenated, and Halogenated Derivatives by Differential Calorimetry
-----	--------	---------	--------	--

hfust	19.40	kJ/mol	354.40	NIST Webbook	
hsubt	110.00 ± 1.20	kJ/mol	329.50	NIST Webbook	
hsubt	99.30 ± 0.20	kJ/mol	308.00	NIST Webbook	
hsubt	98.70 ± 1.70	kJ/mol	316.50	NIST Webbook	

Sources

Crippen Method: http://pubs.acs.org/doi/abs/10.1021/ci990307l

Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws

Heat Capacities of Uracil, Thymine, and https://www.doi.org/10.1021/je060257y

Its Alkylated, Cyclooligomethylenated, McGnarded Maler Derivatives by

http://link.springer.com/article/10.1007/BF02311772

Differential Calorimetry: NIST Webbook:http://webbook.nist.gov/cgi/cbook.cgi?ID=C31703089&Units=SI

Legend

cps: Solid phase heat capacity

hfust: Enthalpy of fusion at a given temperature

hsubt: Enthalpy of sublimation at a given temperature

log10ws: Log10 of Water solubility in mol/llogp: Octanol/Water partition coefficientmcvol: McGowan's characteristic volume

Latest version available from:

https://www.chemeo.com/cid/67-566-0/1-3-Dimethyl-5-ethyluracil.pdf

Generated by Cheméo on 2024-05-11 04:41:48.82175945 +0000 UTC m=+17691757.742336762.

Cheméo (https://www.chemeo.com) is the biggest free database of chemical and physical data for the process industry.