

11«alpha»-Hydroxy-17«alpha»-methyl testosterone

Other names:

4-Androsten-17«alpha»-methyl-11«alpha»,17«beta»-diol-3-one
Androst-4-en-3-one, 11,17-dihydroxy-17-methyl-, (11«alpha»,17«beta»)-
11-Hydroxy-17-methyltestosterone
11Beta-17beta-dihydroxy-17alpha-methylandrost-4-en-3-one
11«alpha»,17«beta»-dihydroxy-17-methylandrost-4-en-3-one

Inchi:

InChI=1S/C20H30O3/c1-18-8-6-13(21)10-12(18)4-5-14-15-7-9-20(3,23)19(15,2)11-16(22)

InchiKey:

UBIBSXMTVJAYTQ-ISQMNYQQSA-N

Formula:

C₂₀H₃₀O₃

SMILES:

CC12CCC(=O)C=C1CCC1C2C(O)CC2(C)C1CCC2(C)O

Mol. weight [g/mol]:

318.45

CAS:

1807-02-9

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -115.48 | kJ/mol | Joback Method |
| hf | -606.88 | kJ/mol | Joback Method |
| hfus | 22.43 | kJ/mol | Joback Method |
| hvap | 94.80 | kJ/mol | Joback Method |
| log10ws | -4.45 | | Crippen Method |
| logp | 3.240 | | Crippen Method |
| mcvol | 258.230 | ml/mol | McGowan Method |
| pc | 2034.55 | kPa | Joback Method |
| tb | 948.34 | K | Joback Method |
| tc | 1178.40 | K | Joback Method |
| tf | 631.44 | K | Joback Method |
| vc | 0.966 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|---------|---------|-----------------|---------------|
| cpg | 984.00 | J/mol×K | 948.34 | Joback Method |
| cpg | 1012.73 | J/mol×K | 986.68 | Joback Method |
| cpg | 1043.17 | J/mol×K | 1025.03 | Joback Method |
| cpg | 1075.71 | J/mol×K | 1063.37 | Joback Method |

| | | | | |
|-----|---------|---------|---------|---------------|
| cpg | 1110.76 | J/mol×K | 1101.71 | Joback Method |
| cpg | 1148.70 | J/mol×K | 1140.05 | Joback Method |
| cpg | 1189.93 | J/mol×K | 1178.40 | Joback Method |

Sources

| | |
|------------------------|---|
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=C1807029&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/80-553-9/11-alpha-Hydroxy-17-alpha-methyl-testosterone.pdf>

Generated by Cheméo on 2025-12-05 09:24:02.306651874 +0000 UTC m=+4674839.836692528.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.