

Glycine, 2-cyclohexyl-N-propoxycarbonyl-, pentyl ester

Inchi: InChI=1S/C17H31NO4/c1-3-5-9-13-21-16(19)15(14-10-7-6-8-11-14)18-17(20)22-12-4-2/
InchiKey: KSZNXRDTZOAJP-UHFFFAOYSA-N
Formula: C17H31NO4
SMILES: CCCCCOC(=O)C(N=C(O)OCCC)C1CCCCC1
Mol. weight [g/mol]: 313.43

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|--------|----------------|
| hf | -801.99 | kJ/mol | Joback Method |
| hvap | 85.12 | kJ/mol | Joback Method |
| log10ws | -4.13 | | Crippen Method |
| logp | 4.009 | | Crippen Method |
| mcvol | 264.390 | ml/mol | McGowan Method |
| pc | 1461.25 | kPa | Joback Method |
| rinpol | 2123.00 | | NIST Webbook |
| tb | 874.92 | K | Joback Method |
| tc | 1079.48 | K | Joback Method |

Sources

Crippen Method: <http://pubs.acs.org/doi/abs/10.1021/ci9903071>
Crippen Method: https://www.chemeo.com/doc/models/crippen_log10ws
Joback Method: https://en.wikipedia.org/wiki/Joback_method
McGowan Method: <http://link.springer.com/article/10.1007/BF02311772>
NIST Webbook: <http://webbook.nist.gov/cgi/cbook.cgi?ID=U383067&Units=SI>

Legend

hf: Enthalpy of formation at standard conditions
hvap: Enthalpy of vaporization at standard conditions
log10ws: Log10 of Water solubility in mol/l

| | |
|----------------|-------------------------------------|
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |

Latest version available from:

<https://www.cheméo.com/cid/83-340-2/Glycine-2-cyclohexyl-N-propoxycarbonyl-pentyl-ester.pdf>

Generated by Cheméo on 2024-04-26 09:22:39.366753161 +0000 UTC m=+16412608.287330473.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.