

Fumaric acid, eicosyl 2-methylpentyl ester

Inchi: InChI=1S/C30H56O4/c1-4-6-7-8-9-10-11-12-13-14-15-16-17-18-19-20-21-22-26-33-29(3)
InchiKey: ZIXMAAYGENNJQA-OCOZRVBESA-N
Formula: C30H56O4
SMILES: CCCCCCCCCCCCCCCCCCCCCOC(=O)C=CC(=O)OCC(C)CCC
Mol. weight [g/mol]: 480.76

Physical Properties

| Property code | Value | Unit | Source |
|---------------|----------|----------------------|----------------|
| gf | -188.34 | kJ/mol | Joback Method |
| hf | -1040.19 | kJ/mol | Joback Method |
| hfus | 75.71 | kJ/mol | Joback Method |
| hvap | 100.26 | kJ/mol | Joback Method |
| log10ws | -9.72 | | Crippen Method |
| logp | 9.107 | | Crippen Method |
| mvol | 444.140 | ml/mol | McGowan Method |
| pc | 642.22 | kPa | Joback Method |
| rinpol | 3284.00 | | NIST Webbook |
| rinpol | 3284.00 | | NIST Webbook |
| tb | 1042.10 | K | Joback Method |
| tc | 1301.69 | K | Joback Method |
| tf | 552.10 | K | Joback Method |
| vc | 1.738 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 1577.07 | J/molxK | 1042.10 | Joback Method |
| cpg | 1674.27 | J/molxK | 1258.42 | Joback Method |
| cpg | 1658.60 | J/molxK | 1215.16 | Joback Method |
| cpg | 1641.18 | J/molxK | 1171.89 | Joback Method |
| cpg | 1621.87 | J/molxK | 1128.63 | Joback Method |
| cpg | 1600.54 | J/molxK | 1085.36 | Joback Method |
| cpg | 1688.31 | J/molxK | 1301.69 | Joback Method |
| dvisc | 0.0000092 | Paxs | 1042.10 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0000126 | Paxs | 960.43 | Joback Method |
| dvisc | 0.0000183 | Paxs | 878.77 | Joback Method |
| dvisc | 0.0000288 | Paxs | 797.10 | Joback Method |
| dvisc | 0.0000502 | Paxs | 715.43 | Joback Method |
| dvisc | 0.0001011 | Paxs | 633.77 | Joback Method |
| dvisc | 0.0002504 | Paxs | 552.10 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U348738&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|----------------------------|---|
| cp_g: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log₁₀ws: | Log ₁₀ of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/88-379-5/Fumaric-acid-eicosyl-2-methylpentyl-ester.pdf>

Generated by Cheméo on 2024-04-25 18:33:52.409382486 +0000 UTC m=+16359281.329959801.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.