

1,2-Cyclohexanedicarboxylic acid, butyl 2-methylpent-3-yl ester

| | |
|----------------------|---|
| Inchi: | InChI=1S/C18H32O4/c1-5-7-12-21-17(19)14-10-8-9-11-15(14)18(20)22-16(6-2)13(3)4/h1 |
| InchiKey: | MJUGEXJVKKJSGY-UHFFFAOYSA-N |
| Formula: | C18H32O4 |
| SMILES: | CCCCOC(=O)C1CCCCC1C(=O)OC(CC)C(C)C |
| Mol. weight [g/mol]: | 312.44 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -355.30 | kJ/mol | Joback Method |
| hf | -881.03 | kJ/mol | Joback Method |
| hfus | 33.81 | kJ/mol | Joback Method |
| hvap | 73.32 | kJ/mol | Joback Method |
| log10ws | -4.36 | | Crippen Method |
| logp | 4.114 | | Crippen Method |
| mcvol | 268.500 | ml/mol | McGowan Method |
| pc | 1406.95 | kPa | Joback Method |
| rinpol | 2075.00 | | NIST Webbook |
| tb | 777.82 | K | Joback Method |
| tc | 976.97 | K | Joback Method |
| tf | 410.08 | K | Joback Method |
| vc | 1.012 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 849.75 | J/molxK | 777.82 | Joback Method |
| cpg | 934.35 | J/molxK | 943.78 | Joback Method |
| cpg | 919.99 | J/molxK | 910.59 | Joback Method |
| cpg | 904.36 | J/molxK | 877.39 | Joback Method |
| cpg | 887.45 | J/molxK | 844.20 | Joback Method |
| cpg | 869.25 | J/molxK | 811.01 | Joback Method |
| cpg | 947.45 | J/molxK | 976.97 | Joback Method |
| dvisc | 0.0000784 | Paxs | 777.82 | Joback Method |
| dvisc | 0.0001054 | Paxs | 716.53 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0001498 | Paxs | 655.24 | Joback Method |
| dvisc | 0.0002290 | Paxs | 593.95 | Joback Method |
| dvisc | 0.0003860 | Paxs | 532.66 | Joback Method |
| dvisc | 0.0007450 | Paxs | 471.37 | Joback Method |
| dvisc | 0.0017505 | Paxs | 410.08 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U339442&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/88-920-3/1-2-Cyclohexanedicarboxylic-acid-butyl-2-methylpent-3-yl-ester.pdf>

Generated by Cheméo on 2024-04-29 11:35:00.12984862 +0000 UTC m=+16679749.050425931.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.