

Succinic acid, 2-ethylhexyl 3-methylpentyl ester

| | |
|----------------------|---|
| Inchi: | InChI=1S/C18H34O4/c1-5-8-9-16(7-3)14-22-18(20)11-10-17(19)21-13-12-15(4)6-2/h15-1 |
| InchiKey: | WGRTWXITQZJMFQ-UHFFFAOYSA-N |
| Formula: | C18H34O4 |
| SMILES: | CCCCC(CC)COC(=O)CCC(=O)OCCC(C)CC |
| Mol. weight [g/mol]: | 314.46 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -372.04 | kJ/mol | Joback Method |
| hf | -915.01 | kJ/mol | Joback Method |
| hfus | 40.90 | kJ/mol | Joback Method |
| hvap | 73.20 | kJ/mol | Joback Method |
| log10ws | -4.60 | | Crippen Method |
| logp | 4.506 | | Crippen Method |
| mvol | 279.360 | ml/mol | McGowan Method |
| pc | 1241.58 | kPa | Joback Method |
| rinpol | 2042.00 | | NIST Webbook |
| rinpol | 2042.00 | | NIST Webbook |
| tb | 762.94 | K | Joback Method |
| tc | 944.42 | K | Joback Method |
| tf | 406.94 | K | Joback Method |
| vc | 1.079 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 850.46 | J/molxK | 762.94 | Joback Method |
| cpg | 868.17 | J/molxK | 793.19 | Joback Method |
| cpg | 884.91 | J/molxK | 823.43 | Joback Method |
| cpg | 900.70 | J/molxK | 853.68 | Joback Method |
| cpg | 915.55 | J/molxK | 883.93 | Joback Method |
| cpg | 929.48 | J/molxK | 914.17 | Joback Method |
| cpg | 942.50 | J/molxK | 944.42 | Joback Method |
| dvisc | 0.0015163 | Paxs | 406.94 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0006283 | Paxs | 466.27 | Joback Method |
| dvisc | 0.0003176 | Paxs | 525.61 | Joback Method |
| dvisc | 0.0001844 | Paxs | 584.94 | Joback Method |
| dvisc | 0.0001183 | Paxs | 644.27 | Joback Method |
| dvisc | 0.0000818 | Paxs | 703.61 | Joback Method |
| dvisc | 0.0000599 | Paxs | 762.94 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U390647&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/89-001-2/Succinic-acid-2-ethylhexyl-3-methylpentyl-ester.pdf>

Generated by Cheméo on 2024-04-28 20:19:16.305485152 +0000 UTC m=+16624805.226062463.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.