

Succinic acid, dec-2-yl 3-methylbut-3-en-1-yl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C19H34O4/c1-5-6-7-8-9-10-11-17(4)23-19(21)13-12-18(20)22-15-14-16(2)3/h |
| InchiKey: | LYMJJXQYBKKJPD-UHFFFAOYSA-N |
| Formula: | C19H34O4 |
| SMILES: | <chem>C=C(C)CCOC(=O)CCC(=O)OC(C)CCCCCCC</chem> |
| Mol. weight [g/mol]: | 326.47 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -281.89 | kJ/mol | Joback Method |
| hf | -814.73 | kJ/mol | Joback Method |
| hfus | 44.43 | kJ/mol | Joback Method |
| hvap | 75.22 | kJ/mol | Joback Method |
| log10ws | -5.47 | | Crippen Method |
| logp | 4.958 | | Crippen Method |
| mvol | 289.150 | ml/mol | McGowan Method |
| pc | 1193.17 | kPa | Joback Method |
| rinpol | 2128.00 | | NIST Webbook |
| rinpol | 2128.00 | | NIST Webbook |
| tb | 782.82 | K | Joback Method |
| tc | 966.60 | K | Joback Method |
| tf | 417.49 | K | Joback Method |
| vc | 1.123 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 882.64 | J/mol×K | 782.82 | Joback Method |
| cpg | 900.14 | J/mol×K | 813.45 | Joback Method |
| cpg | 916.67 | J/mol×K | 844.08 | Joback Method |
| cpg | 932.25 | J/mol×K | 874.71 | Joback Method |
| cpg | 946.88 | J/mol×K | 905.34 | Joback Method |
| cpg | 960.59 | J/mol×K | 935.97 | Joback Method |
| cpg | 973.40 | J/mol×K | 966.60 | Joback Method |

Sources

| | |
|------------------------|---|
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U391141&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307I |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/89-481-0/Succinic-acid-dec-2-yl-3-methylbut-3-en-1-yl-ester.pdf>

Generated by Cheméo on 2024-04-26 04:16:42.331130931 +0000 UTC m=+16394251.251708246.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.