

Succinic acid, di(2-methylbutyl) ester

Inchi: InChI=1S/C14H26O4/c1-5-11(3)9-17-13(15)7-8-14(16)18-10-12(4)6-2/h11-12H,5-10H2,1
InchiKey: ZJVNYQFZJFPWOJ-UHFFFAOYSA-N
Formula: C14H26O4
SMILES: CCC(C)COC(=O)CCC(=O)OCC(C)CC
Mol. weight [g/mol]: 258.35

Physical Properties

Property code	Value	Unit	Source
gf	-405.72	kJ/mol	Joback Method
hf	-832.45	kJ/mol	Joback Method
hfus	30.54	kJ/mol	Joback Method
hvap	64.29	kJ/mol	Joback Method
log10ws	-2.92		Crippen Method
logp	2.945		Crippen Method
mcvol	223.000	ml/mol	McGowan Method
pc	1660.55	kPa	Joback Method
rinpol	1680.00		NIST Webbook
rinpol	1680.00		NIST Webbook
tb	671.42	K	Joback Method
tc	852.27	K	Joback Method
tf	361.86	K	Joback Method
vc	0.856	m3/kmol	Joback Method

Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	623.10	J/mol×K	671.42	Joback Method
cpg	696.05	J/mol×K	822.13	Joback Method
cpg	683.00	J/mol×K	791.99	Joback Method
cpg	669.19	J/mol×K	761.84	Joback Method
cpg	654.60	J/mol×K	731.70	Joback Method
cpg	639.24	J/mol×K	701.56	Joback Method
cpg	708.34	J/mol×K	852.27	Joback Method
dvisc	0.0001031	Pa×s	671.42	Joback Method

dvisc	0.0001395	Pa×s	619.83	Joback Method
dvisc	0.0001994	Pa×s	568.23	Joback Method
dvisc	0.0003062	Pa×s	516.64	Joback Method
dvisc	0.0005173	Pa×s	465.05	Joback Method
dvisc	0.0009958	Pa×s	413.45	Joback Method
dvisc	0.0023105	Pa×s	361.86	Joback Method

Sources

NIST Webbook:	http://webbook.nist.gov/cgi/cbook.cgi?ID=U389643&Units=SI
Crippen Method:	http://pubs.acs.org/doi/abs/10.1021/ci990307l
Crippen Method:	https://www.chemeo.com/doc/models/crippen_log10ws
Joback Method:	https://en.wikipedia.org/wiki/Joback_method
McGowan Method:	http://link.springer.com/article/10.1007/BF02311772

Legend

cp_g:	Ideal gas heat capacity
dvisc:	Dynamic viscosity
g_f:	Standard Gibbs free energy of formation
h_f:	Enthalpy of formation at standard conditions
h_{fus}:	Enthalpy of fusion at standard conditions
h_{vap}:	Enthalpy of vaporization at standard conditions
log_{10ws}:	Log10 of Water solubility in mol/l
log_p:	Octanol/Water partition coefficient
mc_{vol}:	McGowan's characteristic volume
p_c:	Critical Pressure
rin_{pol}:	Non-polar retention indices
t_b:	Normal Boiling Point Temperature
t_c:	Critical Temperature
t_f:	Normal melting (fusion) point
v_c:	Critical Volume

Latest version available from:

<https://www.chemeo.com/cid/89-951-8/Succinic-acid-di-2-methylbutyl-ester.pdf>

Generated by Cheméo on 2025-12-06 01:45:59.886988905 +0000 UTC m=+4733757.417029559.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.