

Succinic acid, 6-methylhept-2-yl 2-methylhex-3-yl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C19H36O4/c1-7-9-17(15(4)5)23-19(21)13-12-18(20)22-16(6)11-8-10-14(2)3/h |
| InchiKey: | ZZVQMNHJGCGEGZ-UHFFFAOYSA-N |
| Formula: | C19H36O4 |
| SMILES: | CCCC(OC(=O)CCC(=O)OC(C)CCCC(C)C)C(C)C |
| Mol. weight [g/mol]: | 328.49 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -368.50 | kJ/mol | Joback Method |
| hf | -946.21 | kJ/mol | Joback Method |
| hfus | 36.45 | kJ/mol | Joback Method |
| hvap | 74.65 | kJ/mol | Joback Method |
| log10ws | -5.24 | | Crippen Method |
| logp | 4.893 | | Crippen Method |
| mvol | 293.450 | ml/mol | McGowan Method |
| pc | 1174.44 | kPa | Joback Method |
| rinpol | 1976.00 | | NIST Webbook |
| rinpol | 1976.00 | | NIST Webbook |
| tb | 784.94 | K | Joback Method |
| tc | 970.70 | K | Joback Method |
| tf | 388.21 | K | Joback Method |
| vc | 1.123 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|-----------|---------|-----------------|---------------|
| cpg | 910.84 | J/molxK | 784.94 | Joback Method |
| cpg | 929.17 | J/molxK | 815.90 | Joback Method |
| cpg | 946.44 | J/molxK | 846.86 | Joback Method |
| cpg | 962.66 | J/molxK | 877.82 | Joback Method |
| cpg | 977.85 | J/molxK | 908.78 | Joback Method |
| cpg | 992.02 | J/molxK | 939.74 | Joback Method |
| cpg | 1005.19 | J/molxK | 970.70 | Joback Method |
| dvisc | 0.0021477 | Paxs | 388.21 | Joback Method |

| | | | | |
|-------|-----------|------|--------|---------------|
| dvisc | 0.0007008 | Paxs | 454.33 | Joback Method |
| dvisc | 0.0003040 | Paxs | 520.45 | Joback Method |
| dvisc | 0.0001591 | Paxs | 586.58 | Joback Method |
| dvisc | 0.0000950 | Paxs | 652.70 | Joback Method |
| dvisc | 0.0000624 | Paxs | 718.82 | Joback Method |
| dvisc | 0.0000439 | Paxs | 784.94 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U381364&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|----------------------------|---|
| cp_g: | Ideal gas heat capacity |
| dvisc: | Dynamic viscosity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log₁₀ws: | Log ₁₀ of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| m_cvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/91-672-5/Succinic-acid-6-methylhept-2-yl-2-methylhex-3-yl-ester.pdf>

Generated by Cheméo on 2024-04-29 01:36:05.527557264 +0000 UTC m=+16643814.448134576.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.