

Hexanoic acid, 2,7-dimethyloct-7-en-5-yn-4-yl ester

| | |
|----------------------|---|
| Inchi: | InChI=1S/C16H26O2/c1-6-7-8-9-16(17)18-15(12-14(4)5)11-10-13(2)3/h14-15H,2,6-9,12H |
| InchiKey: | DBEQUMZHKFWGOD-UHFFFAOYSA-N |
| Formula: | C16H26O2 |
| SMILES: | <chem>C=C(C)C#CC(CC(C)C)OC(=O)CCCC</chem> |
| Mol. weight [g/mol]: | 250.38 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 127.13 | kJ/mol | Joback Method |
| hf | -240.99 | kJ/mol | Joback Method |
| hfus | 33.47 | kJ/mol | Joback Method |
| hvap | 61.15 | kJ/mol | Joback Method |
| log10ws | -4.90 | | Crippen Method |
| logp | 4.104 | | Crippen Method |
| mcvol | 230.840 | ml/mol | McGowan Method |
| pc | 1619.37 | kPa | Joback Method |
| rinpola | 1569.00 | | NIST Webbook |
| rinpola | 1569.00 | | NIST Webbook |
| tb | 646.45 | K | Joback Method |
| tc | 839.85 | K | Joback Method |
| tf | 402.62 | K | Joback Method |
| vc | 0.887 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 612.78 | J/molxK | 646.45 | Joback Method |
| cpg | 630.61 | J/molxK | 678.68 | Joback Method |
| cpg | 647.55 | J/molxK | 710.92 | Joback Method |
| cpg | 663.62 | J/molxK | 743.15 | Joback Method |
| cpg | 678.85 | J/molxK | 775.39 | Joback Method |
| cpg | 693.26 | J/molxK | 807.62 | Joback Method |
| cpg | 706.88 | J/molxK | 839.85 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U299357&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpola: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/92-519-4/Hexanoic-acid-2-7-dimethyloct-7-en-5-yn-4-yl-ester.pdf>

Generated by Cheméo on 2024-04-24 08:34:15.873010189 +0000 UTC m=+16236904.793587511.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.