

Benzamide, 2-fluoro-N-methyl-

| | |
|-----------------------------|---|
| Inchi: | InChI=1S/C8H8FNO/c1-10-8(11)6-4-2-3-5-7(6)9/h2-5H,1H3,(H,10,11) |
| InchiKey: | NAGFMACWWJYORB-UHFFFAOYSA-N |
| Formula: | C8H8FNO |
| SMILES: | CN=C(O)c1ccccc1F |
| Mol. weight [g/mol]: | 153.15 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|--------|----------------|
| hf | -259.30 | kJ/mol | Joback Method |
| hvap | 55.60 | kJ/mol | Joback Method |
| log10ws | -1.69 | | Crippen Method |
| logp | 1.760 | | Crippen Method |
| mcvol | 113.140 | ml/mol | McGowan Method |
| pc | 3368.44 | kPa | Joback Method |
| rinpol | 1354.00 | | NIST Webbook |
| rinpol | 1354.00 | | NIST Webbook |
| tb | 582.11 | K | Joback Method |
| tc | 791.12 | K | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U407127&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307I |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |

Legend

| | |
|--------------|---|
| hf: | Enthalpy of formation at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |

| | |
|-----------------|-------------------------------------|
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |

Latest version available from:

<https://www.cheméo.com/cid/98-756-5/Benzamide-2-fluoro-N-methyl.pdf>

Generated by Cheméo on 2024-05-01 02:52:57.788346551 +0000 UTC m=+16821226.708923866.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.